



FACHBEREICH PHYSIK
BERGISCHE UNIVERSITÄT
GESAMTHOCHSCHULE WUPPERTAL

The Next Round of Hadronic Generator Tuning Heavily Based on Identified Particle Data

Klaus Hamacher and Michael Weierstall

Wuppertal, Juni 1995
WU B 95-07
DELPHI 95-80 PHYS 515
hep-ex/951127



FACHBEREICH PHYSIK
BERGISCHE UNIVERSITÄT
GESAMTHOCHSCHULE WUPPERTAL

The Next Round of Hadronic Generator Tuning Heavily Based on Identified Particle Data

Klaus Hamacher and Michael Weierstall

Abstract

Event shape and charged particle inclusive distributions determined from 750 000 hadronic Z events measured with the DELPHI detector at LEP are presented. The statistical and systematic precision of this data allows for a decisive confrontation with Monte Carlo models of the hadronization process and a better understanding of the structure of the Z hadronic final state.

Improved tunings of the JETSET, ARIADNE and HERWIG parton shower models and the JETSET matrix element model are obtained by fitting the models to identified particle distributions from all LEP experiments and the DELPHI data presented. The description of the data distributions by the models is critically reviewed with special importance attributed to identified particles.

Wuppertal, Juni 1995
WU B 95-07
DELPHI 95-80 PHYS 515

1 Introduction

Precision measurements at LEP using the hadronic final state, such as determinations of the strong coupling constant α_s from event shapes, the measurement of the Z mass and width, the forward backward asymmetries for quarks or at higher energies the $W^\pm$ mass require a precise modelling of the properties of the corresponding final states. Perturbative QCD cannot provide full theoretical insight to the transition of the primary quarks to observable hadrons, the so-called fragmentation or hadronization process. Only part of this transition including large momentum transfer, mainly the radiation of hard gluons or the evolution of a parton shower are calculable perturbatively. The final formation of hadrons is hidden due to the increase of the strong coupling constant α_s at small momentum transfer and hence the failure of perturbation theory.

Guidance to a better understanding of the hadronization process must therefore come from detailed experimental investigations of the hadronic final state including attempts to describe this process by phenomenological models inspired by QCD. LEP I is a unique unrivaled place to pursue these studies. The clean well defined initial state in e^+e^- annihilation provides an excellent testing field, since the event rate at the Z is huge, the energy is large, and the capabilities of the experimental apparatus are much improved with respect to previous experiments.

This paper attempts to determine parameters for the most frequently used hadronization models which lead to an optimal description of the observed hadronic event shapes and charged particle inclusive distributions, as measured with the DELPHI experiment at LEP, as well as the available information on identified particles from all LEP experiments. The latter allows one to precisely determine more model parameters than from event shapes only and also to check the internal consistency of the models. The performance of these models is compared and critically reviewed.

This paper is organized as follows: Section 2 gives a brief overview to the relevant detector components and describes the experimental procedure applied to determine event shape and inclusive distributions and the related systematic errors. Section 3 discusses the models employed and the relevant parameters. Section 4 describes the optimization strategy applied to obtain best parameters for the fragmentation models and justifies the choice of distributions used in the fit. The fits are discussed in detail and the resulting optimized parameters and their errors are presented. In Section 5 a confrontation of the models to the event shape, charge particle spectra and identified particle data is presented. Finally we summarize in Section 6. The appendices contain the definitions of the variables used throughout this paper, tables of model parameter settings and tables and plots of the relevant data distributions and model comparisons.

2 Detector and Data Analysis

The data used in this analysis are from the 1991, 1992 and 1993 data taking with the DELPHI detector at LEP. For the determination of the event shape distributions we use charged particles measured in the solenoidal 1.2 T magnetic field of DELPHI and neutral clusters measured with the electromagnetic or hadronic calorimeters. The following detectors are relevant to the analysis [1]:

- The Vertex Detector VD, measuring the $R\varphi$ coordinate with up to three layers of

silicon micro strips at radii between 6.3 cm and 11 cm. It covers polar angles θ between 37° and 143° .

- The Inner Detector ID, a 24 layer cylindrical jet chamber (θ coverage 17° to 163°).
- The Time Projection Chamber TPC, the principal tracker of DELPHI, with twice 6 sector plates, 24 pad rows and 192 sense wires each. Inner and outer radius of the TPC are 30 cm and 122 cm, the polar coverage is 20° to 160° .
- The Outer Detector OD, a five layer drift chamber at 192 cm radius covering polar angles between 43° and 137° .
- Two sets of forward planar drift chambers FCA and FCB with 6 or 12 layers respectively with a overall polar angle coverage between 11° to 35° and 145° to 169° .
- The High density Projection Chamber HPC, a lead-gas electromagnetic calorimeter with a very good spatial resolution inside the DELPHI coil between 208 cm to 260 cm radius. It measures the showers using the TPC principle and covers polar angles between 43° and 137° . The precision on the energy can be described by $\Delta E/E \approx 29\%/\sqrt{(E)} + 4\%$.
- The Forward Electromagnetic Calorimeter FEMC, a lead-glass array in both end-caps each consisting of 4500 lead glass blocks with a pointing geometry. The angular acceptance is between 10° and 36.5° to the beam. The precision on the energy is $\Delta E/E \approx \sqrt{(0.35 + 5/\sqrt{E})^2 + (6/E)^2} \%$.
- The Hadron Calorimeter HAC, a iron-gas hadronic calorimeter outside the coil consisting of at least 19 layers of streamer tubes and 5 cm thick iron plates also used as flux return. The overall angular coverage is 11.2° to 168.8° . The precision on the energy is $120\%/\sqrt{E}$.

Depending on the polar angle and the detectors included in the track fit the precision on the momentum, p , for charged particles in hadronic events is $\Delta p/p^2 = 0.001$ to 0.01 GeV^{-1} . Charged particles were accepted in the analysis if they satisfy the following criteria:

- $p \geq 200 \text{ MeV}$
- $\Delta p/p \leq 1$
- $20^\circ \leq \theta_{\text{track}} \leq 160^\circ$
- Measured track length ≥ 50 cm.
- Impact parameter with respect to the nominal interaction point within 2 cm perpendicular or 5 cm along the beam.

Furthermore we require that charged particles with large momenta ($p \geq 25 \text{ GeV}$) within the geometrical acceptance of the OD or FCB have indeed been measured by these detectors as well as by the ID or VD. This requirement assures a good momentum resolution for high momentum particles.

Neutral clusters or photons reconstructed from conversions were accepted if:

- $1\text{ GeV} \leq E_{cluster}$ for clusters measured with the HAC,
- $0.5\text{ GeV} \leq E_{cluster/\gamma}$ for electromagnetic clusters or photons.

Because the available statistics is very large, the final experimental error is dominated by the systematic error. Therefore cuts were applied to the event kinematics to assure that the major components of the event were measured in DELPHI with optimal efficiency and resolution, as well as to minimize secondary interactions in the detector material. However care has been taken not too strongly bias the measured distributions by these cuts. Events were selected if they fulfil the following conditions:

- there were at least 5 charged particles selected,
- the total energy of charged particles exceeds 15 GeV and 3 GeV for each half of the detector, defined by the (x,y) plane,
- the polar angle of the thrust axis is between 50° and 85° ,
- the momentum imbalance of the event along the beam direction is $|\sum p_z|/\sqrt{s} \leq 0.15$

In total about 750.000 events satisfy these cuts. The contamination of beam gas events, $\gamma\gamma$ -events, and leptonic events other than $\tau^+\tau^-$, is expected to be less than 0.1% and has been neglected. The influence of $\tau^+\tau^-$ events which have a pronounced 2-jet topology and contain high momentum particles has been determined by a simulation study using events generated by the KORALZ model [2] treated by the full simulation of the DELPHI detector DELSIM [3] and the standard data reconstruction chain. The $\tau^+\tau^-$ contributions have been subtracted from the measured data according to the relative abundance of $\tau^+\tau^-$ ($0.16 \pm 0.03\%$) and hadronic events.

We present differential event shape and inclusive single particle distributions as a function of the physical observables defined in appendix A normalized to the number of hadronic Z decays. Three different types of results are given, cross-sections measured from charged particles only corrected to the initial charged final state, or to the charged plus neutral final state and measured charged plus neutral particles corrected to the full final state.

The observed data distributions were corrected for kinematic cuts, limited acceptance, and resolution of the detector as well as effects due to reinteractions of particles inside the detector material. This correction has been calculated using simulated events generated by JETSET 7.3 [4] treated by the full simulation and analysis chain as described above.

For each bin a correction factor C_i has been calculated as the ratio of the generated and final distributions. Particles with a lifetime bigger than 1 ns were considered as stable in the generated distributions. The correction for initial state photon radiation has been determined separately using events generated by JETSET 7.3 PS with and without initial state radiation as predicted by DYMU3 [5].

The simple unfolding by correction factors in general leads to biases of the final results when the detector smearing is bigger than the bin width used, and when the model does not describe the data well [6]. In our case the model has been tuned to DELPHI data as described in [7] and in this paper and is in good agreement for all distributions considered. The relevant parameter settings are given in appendix B. To keep residual biases small with respect to other systematic errors we ensure that the bin width of all presented distributions is at least as big as the detector resolution.

To account for a possible imperfect representation of the DELPHI detector or secondary processes in the simulation program DELSIM, the cuts given above have been varied over a wide range, including small polar angles for the event axis, demanding events with more than 7 tracks etc. Further cuts have been imposed to exclude the boundaries of the TPC sectors for high momentum particles where the agreement between data and simulation is less good. From the stability of the measured distributions a systematic uncertainty as function of the physical observables has been deduced as the R.M.S. of the deviation with respect to the published central value. As the systematic error is expected to grow proportional to the deviation of the overall correction factor from unity an additional relative systematic uncertainty of 10% of this deviation has been added quadratically to the above value. A further systematic error has been added in quadrature for a few bins where the results of the individual data sets, corresponding to the different years of data-taking were found to be statistically incompatible. This error has been calculated such that the χ^2 per degree of freedom for the merging was 1 when this error was considered in addition. The individual estimates of the systematic uncertainty show similar trends. This a posteriori justifies the methods used, however also indicates correlations between the error estimates which could lead to an overestimate of the systematic uncertainty. The final systematic uncertainty has been smoothened in dependence of the individual variables.

The uncertainties shown in the graphs in Appendix G comparing data and models are final experimental uncertainties adding statistical and systematic uncertainties in quadrature. With the exception of the point-to-point scatter they should be interpreted like a statistical uncertainty. The individual errors are given in the tables in appendix D together with the cross-sections.

3 Monte Carlo Models

In this paper we consider the most frequently used fragmentation models, namely JETSET 7.3/7.4 PS and ME [4], ARIADNE 4.06 [8] and HERWIG 5.8 c [9]. HERWIG and JETSET are complete models describing the parton shower evolution (or matrix element calculation), the hadronization of partons into hadrons, and the subsequent decays of short lived particles. ARIADNE only models the parton shower, the consequent hadronization and decays are treated by the corresponding JETSET routines.

3.1 JETSET 7.3 / 7.4 Parton Shower

JETSET with the parton shower option is used with the settings as given in table 1.

Variable	Value	Variable	Value	Variable	Value
MSTJ(11)	3	MSTJ(12)	3	MSTJ(41)	2
MSTJ(45)	5	MSTJ(46)	3	MSTJ(51)	0
MSTJ(101)	5	MSTJ(107)	0		

Table 1: Setting of JETSET PS Switches

The JETSET parton shower algorithm is a coherent leading log algorithm (LLA) with angular ordering. The shower evolves in the CM frame of the partons obeying energy momentum conservation at each step of the shower. The lowest order 3-jet cross-section is reproduced by rejecting part of the first branchings of the initial $q\bar{q}$ system as predicted by the LLA formalism. Angular ordering of the branchings is explicitly imposed and gluon helicity effects can be included. α_s is running with a scale given by the transverse momentum squared of the branching. The shower evolution is stopped at a mass scale Q_0 , then fragmentation takes over. Q_0 and α_s (i.e. Λ_{QCD}) are parameters of the parton shower part of JETSET.

The fragmentation is performed using the Lund string scheme which can be formulated as an iterative procedure. A string stretches in between the oppositely coloured quark and antiquark via the gluon colour charges. Two close by gluons act similar to a single gluon with equal momentum. Therefore the string model is infrared safe. The longitudinal momentum fraction z of a hadron is determined using the Lund symmetric fragmentation function:

$$f(z) = \frac{(1-z)^a}{z} \cdot \exp -\frac{b \cdot m_t^2}{z}$$

$m_t^2 = m^2 + p_t^2$ is the transverse mass squared of the hadron. a and b are parameters of the fragmentation function. b is universal, a in principle can depend on the quark flavour. For heavy quarks we instead use the Peterson fragmentation function [10]:

$$f(z) = \frac{1}{z \left(1 - \frac{1}{z} - \frac{\epsilon_q}{1-z}\right)^2}$$

with parameters $\epsilon_{b(c)}$ which gives a better description of heavy quark fragmentation. The transverse momenta of hadrons are determined from the p_t of its constituent quarks which in turn is chosen from a tunneling process. This leads to a exponential p_t distribution, thus a Gaussian p_t^2 behaviour. The relevant parameter is the standard deviation of this distribution, σ_q .

The tunneling also determines the quark flavour generated in the string breakup, leading to a dependence $\propto \exp -m_q^2$ and thus negligible heavy quark production in the fragmentation. Already strangeness production is strongly suppressed: $P(u\bar{u}) : P(d\bar{d}) : P(s\bar{s}) = 1 : 1 : \gamma_s \approx 0.3$. Mesons are produced according to their quark contents in the six multiplets with smallest mass, i.e. in the states: 1S_0 , 3S_1 , 1P_1 , 3P_0 , 3P_1 and 3P_2 . Contrary to the formulation implemented in JETSET we define individual production probabilities for these multiplets and light, strange or (both) heavy flavours. Except for the light 3P_0 multiplet the probabilities for the P-multiplets are taken $\propto (1 - P(^1S_0) - P(^3S_1)) \cdot (2s+1)$. In a (2 dimensional) string picture production of particles with angular momentum (i.e. P-states) is suppressed and expected to be small (10% [11]).

Using additional mass relations the tunneling mechanism is also applied to baryon production (replacing a quark by a diquark). Parameters related to baryon production are the relative diquark production rate $P(qq)/P(q)$, an extra strange diquark suppression $P(us)/P(ud)/\gamma_s$ and an extra suppression $P(ud1)/P(ud0)$ of spin 1 diquark relative to spin 0 ones leading to $s = 3/2$ and $s = 1/2$ baryons respectively. Furthermore it turned out to be necessary to include an extra suppression of leading baryons. This extra suppression is not used in heavy quark fragmentation and in the simulation of heavy particle decays by fragmentation because here it leads to soft baryon spectra and a strong overall suppression of baryons.

The parameter related to baryon-meson-baryon production, the so-called Popcorn parameter we leave at its default value, also because so far experimental determinations of this parameter are not fully conclusive (see [27] and references therein). Furthermore we do not include simulation of Bose Einstein interference (by LUBOEI) although with properly chosen parameters this procedure results in good representation of the correlation functions for identical particles as well as in a strongly improved description of light meson resonance lineshapes. The energy momentum rescaling performed in this routine strongly influences angular distributions between particles and multi jet rates. Inclusion of this routine leads to widely different model parameters (compare e.g. the resulting parameters given in section 4 (tables 48,49) and the parameters of the DELPHI simulation (see table 9) which includes Bose Einstein interference).

For JETSET 7.3 and a tuning of ARIADNE 4.06 heavy particles decays have been modified to obtain a better description of the heavy particle branching fractions. The modifications act similarly to those performed in JETSET 7.4.

3.2 JETSET 7.4 Matrix Elements

The parton shower simulation is replaced in the historically older matrix element version of JETSET by the exact second order matrix element calculation which provides up to 4 partons. Two calculations GKS [13] and ERT [14] are available in JETSET. In this paper we consider only the default ERT option because it is expected to be more exact. At PETRA/PEP the predicted 4-jet rate turned out to be too small [15] for a given 3-jet rate. This has been connected with higher order terms missing in the second order calculation. These terms can be partially accounted for by choosing a suitable renormalization scale according to the “optimal perturbation theory” description [16] by choosing a new scale $Q^2 = \mu \cdot s, \mu \leq 1$. To obtain an optimal description we fit the parameter μ together with the Λ_{QCD} parameter and the fragmentation parameters as described in the JETSET PS section 3.1. Formally the scale parameter μ replaces the cutoff parameter Q_0 in the ME fit. JETSET 7.4 ME parameter settings are given in table 2.

Variable	Value	Variable	Value	Variable	Value
MSTJ(11)	3	MSTJ(12)	3	MSTJ(41)	2
MSTJ(45)	5	MSTJ(46)	3	MSTJ(51)	0
MSTJ(101)	2	MSTJ(107)	0	MSTJ(111)	1

Table 2: Setting of JETSET ME Switches

3.3 ARIADNE 4.06

ARIADNE is a particularly elegant formulation of a parton shower based on colour dipoles [12, 8]. The emission of a gluon from a colour dipole, i.e. the initial quark antiquark pair creates two new dipoles, one stretched from the quark to the gluon and one from the gluon to the antiquark. Both in turn can independently radiate further gluons. This ansatz automatically includes ordering in angle (or transverse momentum) as well as azimuthal

dependencies in a proper way. The dipole chain resembles the Lund string picture. Parameters are the QCD scale parameter Λ_{QCD} and the cut off scale p_t^{min} which corresponds to Q_0 for JETSET. The evolution variable is the transverse momentum squared. As for JETSET the first order 3 jet cross section is reproduced in ARIADNE. The ARIADNE running parameters are set as in table 3.

Variable	Value	Variable	Value	Variable	Value
MSTA(1)	1	MSTA(2)	1	MSTA(3)	0
MSTA(5)	1	MSTA(12)	1		
MSTJ(11)	3	MSTJ(12)	3	MSTJ(41)	0
MSTJ(45)	5	MSTJ(46)	3	MSTJ(51)	0
MSTJ(101)	5	MSTJ(105)	0	MSTJ(107)	0

Table 3: Setting of ARIADNE Switches

3.4 HERWIG 5.8c

The evolution of the parton shower in HERWIG is based on the Coherent Parton Branching formalism, an extension of the LLA. It accounts for the leading and sub-leading logarithmic terms arising from soft and (or) collinear gluon emission. HERWIG pays special attention to the simulation of QCD interference phenomena [17]. Most important parameters of the parton shower algorithm are Λ_{QCD} (QCDLAM), the quark masses (RMASS(1-6)) and the effective gluon mass (RMASS(13)). The parton shower in HERWIG 5.8c is matched with the first order 3 jet cross section. At the end of the parton shower evolution gluons are split nonperturbatively into $q\bar{q}$ -pairs.

The hadronization in HERWIG proceeds via the so-called “cluster algorithm” based on the preconfinement characteristic of QCD. The colour charge of a parton is compensated to leading order by a anticolour object which is close by in phase space. Combining colour and anticolour objects low mass clusters with no colour are formed. Higher mass clusters are further split into two lighter ones. The splitting is controlled by the parameters CLMAS and CLPOW. In the decay of a cluster containing a quark from the perturbative phase the direction of this quark is remembered. A gaussian smearing is applied controlled by the parameter CLSMR.

Hadrons then are formed in two body cluster decays according to phase space and spin factors. The particle and hadron transverse momentum are thus dynamically produced as a consequence of the cluster mass spectrum. Particle production in cluster decays is modified by changing the a priori weights for the individual hadron types. These are VECWT, TENWT¹ and DECWT for vector or tensor mesons and decuplet baryons respectively. The a priory weights for quarks are PWT(6).

Light particle decays are simulated in HERWIG using decay tables. Particles including heavy quarks decay via quark decay and subsequent fragmentation.

¹ The 1^{+-} and 0^{++} meson multiplets are not included in HERWIG.

Relevant parameter settings for the tuning of HERWIG are given in table 4.

Variable	Value	Variable	Value	Variable	Value
IPROC	100	SUDORD	1	CLDIR	1

Table 4: Setting of HERWIG parameters

4 Fit of Monte Carlo Models to Experimental Data

Classical optimization strategies like hill climbing methods in general fail to converge if they are directly applied to the optimization of a Monte Carlo model because the physical observables predicted by the models are defined only on a statistical basis (and thus only known within the statistical errors). Moreover, this straight forward strategy requires much computer time and cannot be easily repeated to change the input data to be fitted, or check the influence of systematic errors of the data etc. Therefore, similar to previous work [19, 20, 21] we chose to approximate the dependence of the physical observables on the model parameters analytically. For *each bin of each distribution* we use the quadratic form:

$$MC(\vec{p}_0 + \delta\vec{p})(x) \approx f(\vec{p}_0 + \delta\vec{p})(x) = a_0^{(1)}(x) + \sum_{i=1}^n a_i^{(2)}(x)\delta p_i + \sum_{i,j=1}^n a_{ij}^{(3)}(x)\delta p_i\delta p_j \quad (1)$$

Here $f(x)$ denotes the predicted distribution of a physical observable x , $\vec{p}_0$ is a central parameter setting in the n -dimensional parameter space and δp_i the deviation of parameter i from this setting. The last term includes correlating terms between the model parameters. These terms are not present in a linear approximation as used in [21].

The optimal $m = \frac{n}{2} \cdot (n+3) + 1$ parameters $a^{(k)}$ of the expansion are determined from a fit of ansatz 1 to $l > m$ model reference distributions with different parameter settings. This fit is equivalent to the solution of a system of linear equations:

$$A \cdot \vec{c} = \vec{f} = \vec{MC} \quad (2)$$

where column i of the matrix A contains the parameter variations of model set i :

$$A_{i,1..m} = (1, \delta p_1^{(i)}, \dots, \delta p_n^{(i)}, \delta p_1^{(i)2}, \dots, \delta p_n^{(i)2}, \delta p_1^{(i)} \cdot \delta p_2^{(i)}, \dots, \delta p_n^{(i)} \cdot \delta p_{n-1}^{(i)})$$

$\vec{c}$ is the vector of coefficients $a_{i(j)}^{(k)}(x)$, and $\vec{MC}$ the vector of model predictions corresponding to the parameters $\vec{p}_0 + \delta\vec{p}^{(i)}$. The optimal solution of this, for $l \geq m$ overconstrained linear system is obtained using a standard singular value decomposition method [22, 23].

The parameters of the l reference models (generated with equal statistic) are randomly chosen in parameter space around the central point $\vec{p}_0$. Assuming the (a priori chosen) parameter intervals to be renormalized to ± 1 it turns out to be unimportant, except for minor differences in the statistical precision, whether this volume is a cube or a sphere or when the points are placed on the surface of the sphere. Our choice should ensure that the precision of the fitted linear function is roughly constant within the hypersphere.

Statistical errors of the simulated data sets have been neglected because all sets are generated with fixed statistics, and the variation of the distributions for the different parameter choices is relatively small. Consequently the exact statistical uncertainty of the linear approximation is unknown. However the model statistic is chosen such that the overall statistical error is small compared to the experimental uncertainty of the data.

The optimal parameters p_i , their errors σ_i , and correlation coefficients ϱ_{ij} are then determined from a standard χ^2 -fit using MINUIT [24] of the analytic approximations (1) for *all distributions and bins* considered to the corresponding data. Note that the overall analytic approximation contains $n_{bins} \cdot m$ coefficients $a_{i(j)}^{(k)}(x)$.

If the sensitivity of a distribution to a given model parameter p_i is negligible compared to the sensitivity to other parameters (see also next section), the dependence of the approximation on this parameter has been suppressed by omitting this parameter from the linear system (2). This leads to a better convergence of the minimization and more robust results.

The method has been tested by generating $l + 1 = 51$ simulated sets of 50.000 events each with 6 independent parameters and simultaneously fitting the parameters of one set using the other l sets as reference. Here, the statistical error of the approximation is negligible compared to the statistical error of the test set. The pull distributions $(p_i^{fit} - p_i^{true})/\sigma_i$ are approximately standard normal distributions, i.e. the fitting method is unbiased and produces correct errors.

4.1 Choice of Distributions

The parameters of a fragmentation model have a well-defined physical meaning. However some parameters are directly coupled like a , b , and σ_q in the Lund fragmentation function, or the effect of the parameter on physically observed quantities is obscured by other processes like decays. Therefore the choice of distributions to tune the model (parameters) to is not always evident. So far among the many possible distributions some have been chosen in an ad hoc way, and alternative changes of distributions have been used to estimate systematic errors of parameters [7],[19],[21],[25].

In practice, to keep the influence of statistical errors as small as possible it is clear that the models should be fitted to those distributions which show the strongest dependence on the parameter under consideration and least correlations with others. For each distribution $MC(x)$ we therefore calculated its sensitivity to a given model parameter, i.e. the quantity:

$$S_i(x) = \frac{\delta MC(x)}{MC(x)} \Big|_{p_i} / \frac{\delta p_i}{p_i} \approx \frac{\partial \ln MC(x)}{\partial \ln p_i} \Big|_{p_i}$$

where $\delta MC(x)$ is the change of the distribution MC when changing the model parameter p_i by δp_i around its central value. The fraction $\frac{\delta p_i}{p_i}$ gives all parameters the same normalization.

It is observed, that the sensitivity of single particle inclusive distributions to all model parameters is in general larger than for event shape distributions. Compare for example the sensitivities of inclusive spectra and event shapes to JETSET parameters (figures 1 and 2).

It is seen that almost no sensitivity to the parton shower cut off Q_0 is present in the event shape distributions. For this parameter the x_p -momentum spectrum at large x_p is




Figure 1: Sensitivity single track variables

most important. Correlations among the individual parameters in the inclusive spectra are very strong as can be seen by similar (or opposite) behaviour of the sensitivities for different parameters. Especially the opposite, almost symmetric behaviour of the sensitivities on a and b explains the strong correlation between these parameters and why it is possible to find good descriptions for many different choices of a and b . As has been expected, σ_Q is best determined by the p_t^{out} spectrum or related quantities. The variation of the sensitivities are smallest for the rapidity y . Another important quantity is the charged multiplicity which is known to high precision. Obviously it depends on very many model parameters and details of particle decays.

Event shape distributions measuring the overall shape like T , S or $M_{\text{high}}^2/E_{\text{vis}}^2$ mainly depend on α_s i.e. Λ_{QCD} (see 2) except in the 2-jet region where also fragmentation effects are relevant.

The 3-jet rate as measured by D_2^{Durham} or D_2^{Jade} is best suited to determine Λ_{QCD} because the sensitivities on other parameters is negligible. Thus the Λ_{QCD} -determination can almost be decoupled from the determination of other parameters. This also underlines the reliability of α_s -determinations from jet-rates.

Distributions measuring the aplanarity of events like A , m and also $M_{\text{low}}^2/E_{\text{vis}}^2$ again show increased sensitivities especially on α_s but also strong correlations between the parameters. This explains why quantities measuring differences of event shape variables like O , $M_{\text{diff}}^2/E_{\text{vis}}^2$ and the $AEEC$, contrary to popular belief, depend both on Λ_{QCD} and also on many other fragmentation parameters.

For a comprehensive overview tables 11 and 12 show the sensitivities to JETSET 7.4 and HERWIG 5.8 model parameters averaged geometrically over the bins of the distributions. The averaging dilutes part of the sensitivities (compare e.g. fig. 1 and table 11).

From the above discussion we conclude that for a determination of the general parton-shower and fragmentation parameters the models should be fitted to:

- the inclusive (charged) particle distributions as function of x_p , p_t^{in} and p_t^{out} with







Figure 2: Sensitivity to event shape variables

respect to the thrust or sphericity axis,

- the differential three jet rate D_2 or alternatively R_3 ,
- a combination of event shape distributions like T , M , m , or S , A , P , or M_{high}^2/E_{vis}^2 and M_{low}^2/E_{vis}^2 .

Any combination of event shape distributions or event axes for the p_t -spectra is in principle equivalent and can be used to estimate the stability of the fit and systematic errors of the parameters determined or can be viewed as check of the “predictive power” of the models. However because of the strong correlations among the model parameters correlations need to be considered.

The action of the model parameters linked directly to identified particle production (like the strange quark suppression, γ_s , or the relative probability to form a qq pair to make a baryon, $P(qq)/P(q)$, or others) usually follows the physical interpretation more directly. However identified particle spectra have also high sensitivity to fragmentation parameters (see tables 11 and 12). Therefore the concept described above to determine the optimal dependencies is used.

4.2 Strategy of the Fit

4.2.1 JETSET & ARIADNE

First we perform fits of the general fragmentation parameters (Λ_{QCD} , Q_0 and p_t^{cut} or μ , a , b and σ_q) to the charged particle inclusive and global event shape distributions. For each fit 50 simulated sets with 100.000 events are used. Parameters related to identified particles are set to values similar to those used for the DELPHI simulation (see table 9) which are already known to describe identified particle rates and spectra well.

The average scaled momentum $\langle x_E \rangle$ of charm and beauty particles in the models only depends on $\epsilon_{c(b)}$ and Λ_{QCD} . Stable hadron spectra only weakly depend on heavy quark fragmentation parameters. To further reduce dependencies of stable particle distributions on heavy quark parameters and at the same time assure correct $\langle x_E \rangle$ values for D^* and B mesons we adopt the following procedure: the $\langle x_E \rangle$ values for D^* and B mesons for several choices of Λ_{QCD} and $\epsilon_{c(b)}$ are determined within the model. Then the dependence of $\epsilon_{c(b)}$ on $\langle x_E \rangle_{D^*(B)}$ and Λ_{QCD} is parametrized. This allows to chose for each model set and its given Λ_{QCD} value a corresponding value of $\epsilon_{c(b)}$ such that the model reproduces the average heavy meson scaled momenta as measured by the LEP experiments [26] ($\langle x_E \rangle_{D^*} = 0.504$, $\langle x_E \rangle_B = 0.701$).

The models were then fit to several combinations of event shape and semi-inclusive distributions. The results for Λ_{QCD} and σ_q were relatively stable, the variation for Q_0 or p_t^{cut} was bigger and there were many different solutions for a and b . The latter is due to the strong correlation between these two parameters. Therefore from now on we take one central value of b and treat only a as variable. a is preferred because it is less directly coupled to σ_q in the Lund fragmentation function.

Parameters relevant only to the production of specific particles were then adjusted or fitted to the related data. In overview these are:

- The extra η and η' suppressions were fitted to data from [41],[42].

	fit choice S_i :											
distribution	0	1	2	3	4	5	6	7	8	9	10	11
$\langle N_{ch} \rangle$							•	•	•	•	•	•
x_p	•	•	•	•	•	•	•	•	•	•	•	•
y_T	•	•		•		•	•	•		•		•
y_S	•		•		•	•	•		•		•	•
$p_t^{in}, p_t^{out} \ (T)$	•	•		•		•	•	•		•		•
$p_t^{in}, p_t^{out} \ (S)$	•		•		•	•	•		•		•	•
$S, \ A, \ P$	•		•		•	•	•		•		•	•
$T, \ M, \ m$	•	•		•		•	•	•		•		•
D_2^{Durham}	•	•		•		•	•	•		•		•
D_2^{Jade}	•		•		•	•	•		•		•	•
D_3^{Durham}	•			•		•	•			•		•
D_3^{Jade}	•				•	•	•				•	•
D_4^{Durham}				•		•				•		•
D_4^{Jade}					•	•					•	•

Table 5: Overview of the combinations of inclusive distributions and shape distributions fits of the JETSET, ARIADNE and HERWIG models. Distributions used are marked by •.

- The probability for the different B meson multiplets were adjustet to agree with recent measurements [28], [29]. The corresponding D mesons probabilities were interpolated between the B and light meson values.
- $P(us)/P(ud)/\gamma_s$, The strange baryon suppression was adjusted to the ratio of proton- and Λ^0 -production [27],[30],[33],[32].
- $P(ud1)/P(ud0)$ the spin 1 diquark suppression was fitted to the ratio of $\Sigma(1385)$ and Λ^0 or proton production [27],[31],[30],[32],[33].
- The leading baryon suppression was adjusted to the large momentum tail of the proton and Λ^0 spectra [43],[27],[36].

Then a simultaneous fit of 10 important parameters (see parameters where generated ranges are specified in tables 48 – 52) was prepared generating 100 sets of 100.000 events each. The analytical approximation obtained from these simulated sets was fitted to different choices of event shape and particle distributions. For the event shapes we chose the combinations of variables shown in table 5. The choices for identified particles are shown in table 6. These were used to account for the discrepancies observed in the in the proton data from different experiments and the imperfect representation of the $K^\pm$ spectra in the models.

The Λ^0 data has been excluded from the fit because at small momenta it is not described by the models. Inclusion of this distribution leads to ill defined fits and unstable results. The proton data was only used to determine the diquark suppression parameter $P(qq)/P(q)$ and was excluded from the χ^2 calculation for the fits of the remaining parameters. Baryons at intermediate and large momenta are likely, within the models, to

		fit choice P_i :				
data		0	1	2	3	4
ρ°	DELPHI[39]	•	•	•	•	•
ω	L3[40]	•	•	•	•	•
f_0, f_2	DELPHI[39]	•	•	•	•	•
K^0	ALEPH[27], OPAL[31]	•	•	•	•	•
$K^\pm$	ALEPH[30], DELPHI[33], OPAL[32]	•	•	•		
K^{*0}	OPAL[34]	•	•	•	•	•
$K^{*\pm}$	ALEPH[35], DELPHI[39], OPAL[34]	•	•	•	•	•
Φ	DELPHI[33], OPAL[32]	•	•	•	•	•
p	ALEPH[30], DELPHI[33]	•	•		•	
p	OPAL[32]	•		•		•

Table 6: Overview of the combinations of identified particle data used for the JETSET and ARIADNE fits. Data used are marked by •.

be primary particles. Therefore these data strongly influence the fragmentation function and the related parameters. Since proton and Λ^0 spectra appear poorly described by the models and because it is even necessary to modify the fragmentation function by an extra suppression at large momenta this strong impact on the fit results is felt to be unphysical and therefore has been excluded.

Separate fits were performed for all combinations of choices for shapes and identified particles. To check the stability of the fits the optimization has been started with 6 random start values of fragmentation parameters. The fits were stable and converged to the same solution for about 95% of the cases. Differences mainly showed up for two strongly correlated parameters. For those fits which show more than 1 solution we used the one with the better χ^2 . A full (MINOS [24]) error estimate for all parameters was performed.

The central result corresponds to the combination S6/P0. This combination has been chosen because it contains event shape distributions linear and quadratic in the particle momenta, the charged multiplicity and all relevant identified particle information and thus should result in a comprehensive overall description. Results are given with the statistical and systematic errors in tables 48 – 52. The results for the individual parameters are strongly correlated. The correlation coefficients for the JETSET 7.4 (default decays) and ARIADNE 4.06 (DELPHI decays) are given representatively in appendix F. Beside these statistical correlations further correlations exist due to the different possible choices of input data. The systematic error is the R.M.S obtained from all combinations of input data which turn out to give parameters smaller respectively bigger than the central fit.

The following observations were made comparing results of the different input data choices for the JETSET PS fits:

As should be expected the results for Λ_{QCD} , Q_0 and σ_q are almost independent of the choice of identified particle data. The resulting value of Λ_{QCD} does not depend on the algorithm used (JADE or DURHAM) if only D_2 is included in the fit however if also the higher jet rates were used the Λ_{QCD} value turns out to be somewhat bigger ($\approx 8\%$) in case of the JADE algorithm. Values of Λ_{QCD} and Q_0 are positively correlated. This

implies that the number of final partons is more stable within the models than might be expected from the error of Q_0 alone. Λ_{QCD} , and a and σ_q are anticorrelated. This is due to the a compensation of transverse momentum generated in the parton shower and fragmentation phase of the model.

The results for γ_s are higher if $K^\pm$ and K^0 data are included compared to K^0 only. $P(qq)/P(q)$ is larger if the ALEPH proton spectrum is included. Production parameters for strange vector and pseudoscalar mesons are anticorrelated. If the charged multiplicity is not included and therefore the predicted multiplicity is smaller than the measured result, the primary production probabilities for light pseudoscalar and vector mesons are $P(^1S_0)_{ud} \approx 0.40$ and $P(^3S_1)_{ud} \approx 0.26$ respectively. If the multiplicity is fixed to $N_{ch} \approx 20.9$ these values are 0.28 and 0.29. This implies substantial production probabilities for light p-wave mesons of 0.36 to 0.43.

Most of these observations similar also apply in case of ARIADNE. However here the different choices of D_2, D_3 etc. lead to stable results for Λ_{QCD} . There is a tendency to obtain slightly bigger values from the JADE algorithm (0.245) then from Durham (0.237). This goes again conform with bigger values for p_t^{QCD} from JADE (≈ 0.9) then from DURHAM (≈ 0.6).

For JETSET ME Λ_{QCD} is anticorrelated with the scale μ . The fragmentation parameters have different values as in the PS case. Especially $\sigma_q \approx 0.48$ is much bigger. This partially compensates missing higher orders in the ME ansatz. The values of γ_s and $P(qq)/P(q)$ are smaller compared to the PS case and show a decreased dependence on the choice of the related input data. The probability for p-wave states is much bigger ≈ 0.36 already for the fits with free multiplicity.

4.2.2 HERWIG

HERWIG employs much less parameters than JETSET especially in the hadronization sector of the model. Therefore a simultaneous fit of all model parameters which are found to be important (compare table 12) is easily performed. These parameters are:

- the QCD scale parameter QCDLAM and the gluon mass RMASS(13) as major parameters of the PS phase,
- the cluster fragmentation parameters CLPOW, CLMAX and CLSMR (CLDIR=1) and
- the a priori weights DECWT for decuplet baryons, PWT(3) for strange quarks and PWT(7) for diquarks.

Beside the particle spectra the cluster parameters also strongly influence the stable charged particle and event shape distributions. Fits have therefore been performed to combinations of shape distributions (see table 5) and identified particle data (see table 7). Of special importance is the simultaneous usage of proton and Λ^0 data. The average scaled momenta of heavy mesons are important for the determination of CLPOW and CLMAX.

The different choices of the identified particle data (see table 7) are made such that systematic differences in the data are reflected in the systematic errors of the parameters. The central fit result for HERWIG corresponds to the combination S_6P_5 .

		fit choice P_i :									
data		0	1	2	3	4	5	6	7	8	9
ρ^0	DELPHI[39]	•	•	•	•	•	•	•	•	•	•
ω	L3[40]	•	•	•	•	•	•	•	•	•	•
f_2	DELPHI[39]	•	•	•	•	•	•	•	•	•	•
K^0	ALEPH[27], OPAL[31]	•	•	•	•	•	•	•	•	•	•
$K^\pm$	ALEPH[30], DELPHI[33], OPAL[32]	•	•	•			•	•	•		
K^{*0}	OPAL[32]	•	•	•	•	•	•	•	•	•	•
$K^{*\pm}$	ALEPH[35], DELPHI[39], OPAL[34]	•	•	•	•	•	•	•	•	•	•
Φ	DELPHI[33], OPAL[32]	•	•	•	•	•	•	•	•	•	•
Λ^0	ALEPH[27], DELPHI[43]	•	•		•		•	•		•	
Λ^0	OPAL[34]	•		•		•	•		•		•
p	ALEPH[30], DELPHI[33]	•	•		•		•	•		•	
p	OPAL[32]	•		•		•	•		•		•
η	ALEPH[42]	•	•	•	•	•	•	•	•	•	•
η'	ALEPH[42]	•	•	•	•	•	•	•	•	•	•
$\langle x_E \rangle D^{*\pm}, D^{*0}$	[26]	•	•	•	•	•	•	•	•	•	•
$\langle x_E \rangle B^0, B^\pm$	[26]	•	•	•	•	•	•	•	•	•	•
$\Sigma^\pm(1385)$	DELPHI[38], OPAL [34]	•	•	•	•	•					
Ξ^-	DELPHI[38], OPAL [34]	•	•	•	•	•					
$\Xi_0(1530)$	DELPHI[38], OPAL [34]	•	•	•	•	•					

Table 7: Overview of the combinations of identified particle data used for the HERWIG fits. Data used are marked by •.

Following observations are made comparing results of the different input data choices for the HERWIG fits:

Values obtained for QCDLAM are in general bigger (≈ 0.01) when fitting DURHAM compared to JADE jet rates. Contrary to the JETSET case QCDLAM is smaller (by ≈ 0.005) when differential 2-, 3-, and 4-jet rates are fitted compared to 2-jet only. QCDLAM is insensitive to the inclusion of the charged multiplicity in the fit. The gluon mass RMAS(13) shows some dependency on the selected identified particle information and on the multiplicity. The cluster parameters CLMAX and CLPOW depend on the identified particle spectra only in case that the multiplicity is not included in the fit. CLPOW is higher for JADE than for DURHAM jet rates. CLSMR depends only on the charged particle and event shape information. The inclusion of the baryon decuplet data tends to spoil the description of the octet sector.

5 Comparison of Models to Data

The fits of the individual fragmentation models are compared to DELPHI data in figures 3 - 30 shown in appendix G. The lower insets of the plots depict the relative deviation of the models to the data. Also shown as shaded area in these insets is the total experimental error obtained by adding quadratically the systematic and statistical error in each bin. Except for the point-to-point scatter the error should be interpreted like a statistical " 1σ " uncertainty. For most shape distributions we show on the left side the data as measured from charged particles only, on the right measured from charged and neutral particles. The distributions are corrected to the corresponding final states.

We restrict to comparisons to the following models:

- JETSET 7.3 DELPHI decays labeled JT 7.3 PS
- JETSET 7.4 default decays labeled JT 7.4 PS
- ARIADNE 4.06 DELPHI decays labeled AR 4.06
- HERWIG 5.8 c default decays labeled H 5.8C
- JETSET 7.4 ME default decays labeled JT 7.4 ME

The different decay treatments lead only to negligible differences for the stable charged particle and event shape distributions.

Inclusive Charged Particle Spectra

All models describe well the general trends of the data. Almost no discrepancies show up from a direct data model comparison. More quantitatively the comparison of model and data (lower insets) shows:

The x_p -spectrum (fig. 3) for $x_p < 0.4$ is almost perfectly described by the ARIADNE and JETSET PS models. At large x_p these models slightly underestimate the data. This trend is reduced if the multiplicity is left free in the fit. HERWIG and JETSET ME exhibit a slight wave structure when compared to the data. This structure is also reflected in the rapidity distributions (see fig. 4).

Although the p_t^{in} -distribution (fig. 5) is in agreement to data for all models the tail of the p_t^{out} -distribution (fig. 6) ($p_t^{out} > 1$) disagrees with the data for all models. This tail is almost unaffected by fragmentation but is sensitive to the partonic part of the models. For the ME model a discrepancy might be expected because of missing higher order terms in the second order matrix element calculation. The failure of the parton shower models can possibly be traced back to missing large angle terms in the LLA basic to the models. Accepting these reasons for the failure of the models it should be expected that a matching of the second order calculation and the LLA formalism should lead to an improved description of p_t^{out} and related distributions.

Shape Distributions

The general shape distributions $1 - T$, S , C and B_{sum} (figs. 9,19,25,29) measured from charged particles are described within the small experimental errors (typically 2-3%) by all parton shower models. HERWIG for large values of these observables tends to lie slightly above the data. The agreement to the corresponding charged plus neutral distributions is still satisfactory, although these distributions have not been used in the fit. This illustrates further the quality of the models and of the data. The description of distributions sensitive to transverse momenta in the event plane like M , B_{max} or M_h^2/E_{vis}^2

	JETSET 7.3 PS	JETSET 7.4 PS	ARIADNE 4.06	JETSET 7.4 ME	HERWIG 5.8 C	LEP[44],[46],[47],[49]
Charged Particles						
$\langle N_{ch} \rangle$	20.87	20.81	20.80	20.86	20.94	20.95 $\pm$ 0.21
Pseudoscalar Mesons						
$\pi^\pm$	17.19	17.09	17.13	17.36	17.66	17.1 $\pm$ 0.4
π^0	9.85	9.83	9.82	10.03	9.81	9.9 $\pm$ 0.08
$K^\pm$	2.20	2.23	2.19	2.15	2.11	2.42 $\pm$ 0.13
K^0	2.13	2.17	2.12	2.10	2.08	2.12 $\pm$ 0.06
η	1.07	1.10	1.09	1.16	1.02	0.73 $\pm$ 0.07
$\eta'(958)$	0.10	0.09	0.10	0.10	0.14	0.17 $\pm$ 0.05
D^+	0.19	0.20	0.20	0.20	0.24	0.20 $\pm$ 0.03
D^0	0.46	0.49	0.48	0.49	0.53	0.40 $\pm$ 0.06
$B^\pm, B^0$	0.36	0.36	0.36	0.36	0.36	0.34 $\pm$ 0.06
Scalar Mesons						
$f_0(980)$	0.17	0.16	0.17	0.16		0.14 $\pm$ 0.06
Vector Mesons						
$\rho^0(770)$	1.29	1.27	1.26	1.29	1.43	1.40 $\pm$ 0.1
$K^{*\pm}(892)$	0.78	0.77	0.79	0.77	0.74	0.78 $\pm$ 0.08
$K^{*0}(892)$	0.80	0.77	0.81	0.78	0.74	0.77 $\pm$ 0.09
$\phi(1020)$	0.109	0.107	0.107	0.102	0.099	0.086 $\pm$ 0.018
$D^{*\pm}(2010)$	0.18	0.22	0.19	0.22	0.22	0.17 $\pm$ 0.02
$D^{*0}(2007)$	0.20	0.22	0.20	0.22	0.23	.
Tensor Mesons						
$f_2(1270)$	0.29	0.29	0.29	0.30	0.26	0.31 $\pm$ 0.12
Baryons						
p	0.97	0.97	0.96	0.90	0.78	0.92 $\pm$ 0.11
Λ^0	0.361	0.349	0.365	0.309	0.368	0.348 $\pm$ 0.013
Ξ^-	0.0288	0.0300	0.0300	0.0256	0.0493	0.0238 $\pm$ 0.0024
$\Delta^{++}(1232)$	0.158	0.160	0.136	0.158	0.154	0.077 $\pm$ 0.018
$\Sigma^\pm(1385)$	0.037	0.036	0.032	0.033	0.065	0.0380 $\pm$ 0.0062
$\Xi^0(1530)$	0.0073	0.0069	0.0063	0.0060	0.0249	0.0063 $\pm$ 0.0014
Ω^-	0.0013	0.0019	0.0021	0.0010	0.0077	0.0051 $\pm$ 0.0013
Λ_b^0	0.032	0.033	0.032	0.029	0.007	0.031 $\pm$ 0.016

Table 8: The Production Rates for the different Generators compared to LEP data

(figs. 10,27,23) is only slightly less good (typically better than 5%). Turning to distributions sensitive to transverse momenta out of the event plane m , O , A , M_t^2/E_{vis}^2 , or B_{min} (figs. 11,12,20,24, 28) the following pattern is observed: ARIADNE generally describes the data well, HERWIG tends to overestimate the distributions and JETSET PS is usually below the data for higher values of the observables. The latter is to be expected from the underestimation of the p_t^{out} -distribution by all PS models. A similar pattern is also observed for the jet rates. The differential 2-jet rate D_2 (figs. 13,14) is well described by all models. ARIADNE describes well also the higher jet rates (figs. 15–18), HERWIG overestimates and JETSET PS underestimates them. This behaviour is consistently observed for the JADE as well as for the DURHAM algorithm and for the charged and charged plus neutral data.

The description of the shape distributions by the JETSET ME model is less perfect than for the PS models. There is a shortcome for the extreme 2 jet region, the multijet rates and observables sensitive to radiation out of the event plane. In general however the description by the JETSET ME is quite satisfactory.

Identified Particle Rates

Table 8 compares the particle rates predicted from the models with the current measured LEP averages. After inclusion of the charged multiplicity in the fits it is well described by the models. Neglecting this constraint ARIADNE and JETSET PS predict the multiplicity too low (20.2 – 20.4) and JETSET ME too high (22.7). HERWIG is correct

without constraint.

All meson rates with the exception of the $K^\pm$ and the η rate are well described within errors (2σ). The measured η momentum spectrum is however well described. We therefore suspect a too small error quoted in the extrapolation needed to determine the total η rate. The $K^\pm$ rate is sensitive to heavy quark decays (see below). The octet baryons also agree reasonably. Only HERWIG overestimates the Ξ^- rate by about a factor 2. Some discrepancies show up in the decuplet baryon sector.

Meson Momentum Spectra

In figures 31,32 we compare the models to the identified kaon spectra as function of $\xi_p = \log \frac{1}{x_p}$ from different LEP experiments. For a more quantitative comparison which also allows to judge the agreement among the different experiments we show on the right the relative deviation between the individual data sets and each model as lines. In the fragmentation region at large ξ_p (i.e. small x_p) where the $K^\pm$ data is more precise and therefore dominates the fit the K^0 's are overestimated by $\approx 10 - 20\%$ by all models. At more central momenta all kaons are well described. $K^\pm$ are underestimated in the range ($0.8 < \xi_p < 2.5$). K's from heavy particle decays tend to contribute mainly in this momentum range.

Wrong branching fractions of b-hadrons in the models presumably cause this discrepancy. Indeed a recent DELPHI measurement of the inclusive particle production in b-events supports this interpretation [48]. The difference of $K^\pm$ and K^0 production was found to be 0.60 ± 0.50 . The corresponding value in the models is only ≈ 0.30 for JETSET and ARIADNE and 0.04 for HERWIG. This difference causes the relatively large systematic error on γ_s . An improvement is possible as soon as K measurements for identified b and light quark events become available. Finally at very small ξ_p (i.e. large x_p) there is an indication that the models underestimate slightly the K production.

Fragmentation functions of vector mesons, which are likely to be primary particles, are compared to in figs. 33,34,35. Within the large errors of these resonance measurements all models describe the spectra very well. There is only a tendency to predict a somewhat harder fragmentation than measured. This is most evident for the $\Phi(1020)$ spectrum.

It is important to note that the strange pseudoscalar and vector meson data shown already imply large production rates of higher mass resonances with strangeness. The large ξ_p part of the $K^{0,\pm}$ spectra can only be described if large resonance rates are assumed in the models. Due to the stronger boost (because of their high mass) kaons from resonance decays acquire in average higher momenta (i.e. smaller ξ_p). As the strange vector meson rates are relatively moderate higher mass strange resonances must therefore be present. This reflects in high production probabilities, of the order of 25%, for p-wave resonances with strangeness in JETSET and ARIADNE. HERWIG dynamically predicts similar rates.

Fig. 36 depicts the the good agreement of the model predictions to the measurements of the $f_0(980)$ scalar and $f_2(1270)$ tensor meson resonance. With large production probabilities for p-wave resonances all models describe the measured data well. Note that high production of p-wave states ($> 10\%$) is not expected in the string picture [11].

Baryon Momentum Spectra

The proton spectra of OPAL and ALEPH show a severe discrepancy at high momentum (or small ξ_p) (see fig. 37). The central model fits interpolate between these experimental results. Including only the ALEPH or OPAL data in the fit it turns out that the OPAL

proton spectrum cannot be described by JETSET PS and ARIADNE. The ALEPH data can be reasonably well described. On the other hand the JETSET ME model gives a fair description of the OPAL proton data. The rate predicted by HERWIG is too high at small ξ_p and too low at small ξ_p . At small ξ_p the behaviour of JETSET PS and ARIADNE is similar to HERWIG if the extra baryon suppression is not used. The need for an extra suppression of baryon production may also indicate the presence of nucleon resonances. Although the mass splitting between angular excited baryons and the related ground states is smaller than in the mesons case these states are so far not foreseen in JETSET or HERWIG. Decays of these states would lead to a softening of the spectra of stable baryons and would also modify the predicted correlations between baryon antibaryon pairs.

The experimental situation for the Λ^0 spectrum also shows some discrepancies (see fig. 38). JETSET PS and ARIADNE describe or slightly overestimate the small ξ_p region. HERWIG here is again too high. At large ξ_p where the measurements agree all models underestimate the Λ^0 production by $\approx 15\%$. This region is of special interest because here particle production is expected to be influenced by coherence phenomena.

Fig. 39 shows in a comprehensive overview a comparison of the octet and decuplet baryon production and the model predictions. The discrepancies discussed in the last paragraphs are hidden here due to the large scales. JETSET and ARIADNE describe the gross features of the octet and decuplet baryon production well. The Δ^{++} rate [49] is however overestimated by about a factor 2. HERWIG predicts too hard baryon fragmentation. The relative production rates of the different multiplet states are less well predicted by HERWIG.

6 Summary

From 750.000 $e^+e^- \rightarrow Z \rightarrow \text{hadrons}$ events measured by the DELPHI experiment precise fully corrected semi-inclusive charged particle and event shape distributions have been determined.

A systematic, quantitative study has been undertaken to determine the optimal choice of distributions to tune fragmentation models to. Semi-inclusive charged particle and identified particle distributions constrain the hadronization part of the models whereas 3-jet rate and most event shape distributions mainly control the parton shower parameters (especially Λ_{QCD}) in the models.

Optimal parameters for the ARIADNE 4.06, HERWIG 5.8c and JETSET 7.3 and 7.4 parton shower models and for the JETSET 7.4 matrix element model have been determined. The models were fitted to the measured event properties and inclusive data and to identified particle data measured by the LEP experiments. The fit algorithm employed allowed for a simultaneous fit of up to 10 model parameters. Statistical and systematic errors as well as correlations of the model parameters have been determined.

All models reasonably describe the inclusive and the event shape distributions. The data measured form charged particles and charged plus neutral particles when compared with the corresponding model predictions yield consistent results.

All models underestimate the tail of the p_t^{out} distribution by more than 25%. With this exception the best overall description of event shapes is provided by the ARIADNE 4.06 model. HERWIG 5.8c tends to overestimate and JETSET 7.3/7.4 to underestimate the

production of 4 and more jet events. Correspondingly the tails of shape distributions sensitive to particle production out of the event plane are overestimated (underestimated) by HERWIG (JETSET). The matrix element model JETSET 7.4 ME with optimized scale also provides reasonable predictions. It however shows the expected discrepancies due to missing higher orders in the extreme 2-jet and multijet regions.

Identified meson spectra are fairly well described by all models. It has been found that strong production of p-wave resonances (25 – 40%) has to be considered. This is not expected in a string fragmentation picture.

The gross features of baryon production are described by JETSET and ARIADNE. Some discrepancies however show up in the momentum spectra and in the relative abundance of the individual multiplet states. HERWIG shows stronger discrepancies, especially the predicted fragmentation functions are too hard. In JETSET and ARIADNE a similar tendency has been corrected by an extra leading baryon suppression.

Acknowledgement

We would like to thank L. Lönnblad, I. Knowles, M. Seymour and T. Sjöstrand for useful discussions.

A Definition of Variables

Throughout this paper we use the following definitions of event shape and inclusive particle variables:

A.1 Inclusive Single Particle Variables

Scaled Momentum, x_p

The Scaled Momentum x_p is the absolute momentum $|\vec{p}|$ of a particle scaled to the beam momentum.

Transverse Momenta, p_t^{in}, p_t^{out}

We distinguish the transverse momentum of a particle in the event plane $p_t^{in} = \vec{p} \cdot \vec{n}_{Major}$ and out of the event plane $p_t^{out} = \vec{p} \cdot \vec{n}_{Minor}$. Alternatively we use the axes as defined by the eigenvectors of the quadratic momentum tensor.

Rapidity y

The rapidity is given by:

$$y = \frac{1}{2} \cdot \log \frac{E - p_{\parallel}}{E + p_{\parallel}}$$

where $p_{\parallel}$ is a particles momentum parallel to $\vec{n}_{Thrust}$ or $\vec{n}_{Sphericity}$.

A.2 Event Shapes Variables

Thrust T , Major M , Minor m , Oblateness O

Thrust T [50] and Thrust-axis are defined by:

$$T = \max_{\vec{n}_{Thrust}} \frac{\sum_{i=1}^{N_{particle}} |\vec{p}_i \cdot \vec{n}_{Thrust}|}{\sum_{i=1}^{N_{particle}} |\vec{p}_i|}$$

$\vec{p}$ is the 3-momentum of a particle, $\vec{n}_{Thrust}$ is a unit-vector along the Thrust-axis. Major and Minor are defined via the same expression, however replacing $\vec{n}_{Thrust}$ by $\vec{n}_{Major} \perp \vec{n}_{Thrust}$ or $\vec{n}_{Minor} = \vec{n}_{Major} \times \vec{n}_{Thrust}$ respectively. The oblateness is $O = M - m$.

Sphericity S , Aplanarity A , Planarity P

Ordering the eigenvalues λ of the quadratic momentum tensor:

$$M^{\alpha\beta} = \sum_{i=1}^{N_{particle}} p_i^{\alpha} p_i^{\beta} \quad (\alpha, \beta = 1, 2, 3)$$

$$\lambda_1 \geq \lambda_2 \geq \lambda_3 \quad \lambda_1 + \lambda_2 + \lambda_3 = 1$$

The Sphericity is $S = \frac{3}{2}(\lambda_2 + \lambda_3)$, the Aplanarity $A = \frac{3}{2}\lambda_3$ and the Planarity is $P = \frac{2}{3}(S - 2A)$ [51]. The Sphericity-axis is parallel to the eigenvector corresponding to λ_1 . As the momenta enter quadratically the Sphericity-axis is influenced more strongly by large momentum particles than the Thrust-axis.

C- and D-Parameter

C- and D-Parameter are defined through the eigenvalues λ of the linear momentum tensor [52]:

$$\Theta^{\alpha\beta} = \frac{1}{\sum_{k=1}^{N_{particle}} |\vec{p}_k|} \cdot \sum_{i=1}^{N_{particle}} \frac{p_k^i p_k^j}{|\vec{p}_k|}$$

$$C = 3 \cdot (\lambda_1 \lambda_2 + \lambda_2 \lambda_3 + \lambda_3 \lambda_1) \quad D = 27 \cdot \lambda_1 \lambda_2 \lambda_3$$

Jet Masses $M_{high}^2/E_{vis}^2, M_{low}^2/E_{vis}^2, M_{diff}^2/E_{vis}^2$

Particles are ordered in the two hemispheres of an event separated by the plane normal to $\vec{n}_{Thrust}$.

$$\frac{M_{high}^2}{E_{vis}^2} = \frac{1}{E_{vis}^2} \cdot \max \left(\left(\sum_{\vec{p}_k \cdot \vec{n}_{Thrust} > 0} p_k \right)^2, \left(\sum_{\vec{p}_k \cdot \vec{n}_{Thrust} < 0} p_k \right)^2 \right)$$

For $\frac{M_{low}^2}{E_{vis}^2}$ the maximum in the above formula is replaced by the minimum and $\frac{M_{diff}^2}{E_{vis}^2} = \frac{M_{high}^2}{E_{vis}^2} - \frac{M_{low}^2}{E_{vis}^2}$. This definition differs from the original one by Clavelli [53] but is easier to calculate and therefore preferred by many experiments.

Jet Broadening $B_{max}, B_{min}, B_{sum}, B_{diff}$

Similarly to the definition of the jetmasses the transverse momenta with respect to the Thrust-axis are added up [57], i.e.:

$$B_{\pm} = \frac{\sum_{\pm \vec{p}_i \cdot \vec{n}_{Thrust} > 0} |\vec{p}_i \times \vec{n}_{Thrust}|}{2 \sum_i |\vec{p}_i|}$$

Then:

$$B_{max} = \max B_+, B_- \quad B_{min} = \min B_+, B_- \quad B_{sum} = B_+ + B_- \quad B_{diff} = |B_+ - B_-|$$

Differential Jet Rates $D_i(y)$

Jets are reconstructed using cluster finding algorithms of the JADE typ [54]. For each event and each pair of particles i and j the scaled invariant mass or transverse momentum y_{ij} for the JADE or Durham algorithm respectively are evaluated:

$$y_{ij}^{Jade} = 2E_i E_j / E_{vis}^2 \cdot (1 - \cos \theta_{ij})$$

$$y_{ij}^{Durham} = 2 \min(E_i^2, E_j^2) / E_{vis}^2 \cdot (1 - \cos \theta_{ij})$$

where E_i, E_j are the energies and θ_{ij} the angle between the momentum vectors of the two particles. The particle pair with the lowest value y_{ij} is selected and replaced by a pseudo-particle with four momentum $(p_i + p_j)$, hereby reducing the multiplicity by one. In successive steps the procedure is repeated until the scaled invariant masses of all pairs of (pseudo-)particles are larger than a given resolution

y . The remaining (pseudo-)particles are called jets. The differential 2-jet rate D_2 is derived from the 2-jet rate $R_2 = \frac{N_{2-jets}}{N}$ [55]:

$$D_2(y) = \frac{R_2(y + \Delta y) - R_2(y)}{\Delta y}$$

Higher differential rates follow from the recursion:

$$D_n(y) = \frac{R_n(y + \Delta y) - R_n(y)}{\Delta y} + D_{n-1}(y)$$

Energy-Energy-Correlation EEC and Asymmetry $AEEC$

The Energy-Energy-Correlation EEC is the histogram of angles χ_{ij} between all particles weighted by their scaled energies [56]:

$$EEC(\cos\chi) = \frac{1}{N} \frac{1}{\Delta \cos\chi} \cdot \sum_{events}^N \sum_{i,j} \frac{E_i}{E_{vis}} \frac{E_j}{E_{vis}} \Theta(\Delta \cos\chi - |\cos\chi - \cos\chi_{ij}|)$$

$\cos\chi$ and $\Delta \cos\chi$ are the low edge and width of a bin and Θ is the step function. For $\cos\chi \geq 0$ the asymmetry $AEEC$ of the EEC then is:

$$AEEC(\cos\chi) = EEC(-\cos\chi) - EEC(\cos\chi)$$

B Parameter Settings of the DELPHI Monte Carlo

The DELPHI tuning of JETSET has been obtained by tuning the model to charged particle data from the 1991 and 1992 data taking. Care has been taken to describe especially well the observables relevant to standard precision analyses which are the charged multiplicity, the momentum spectrum, 2-jet rate and Thrust and Sphericity distribution. Also included is the simulation of Bose Einstein interference (by LUBOEI) to obtain a correct description of two particle correlations and light resonance line shapes. The BE parameters are taken from the DELPHI measurement [45] Particle spectra have been adjusted and partially fitted to available data [46]. Heavy particle decays have been adjusted. When using adjusted decay tables this is quoted as DELPHI decays.

Variable	Dec. 93	Sept. 94	Variable	Dec. 93	Sept. 94
MSTJ(11)	3	3	PARJ(81)	0.297	0.346
MSTJ(12)	3	3	PARJ(82)	1.732	2.25
MSTJ(41)	2	2	PARJ(92)	1	1
MSTJ(45)	5	5	PARJ(93)	0.394	0.394
MSTJ(46)	3	3	$P(^1S_0)_{ud}$		.423
MSTJ(51)	2	2	$P(^3S_1)_{ud}$		.275
MSTJ(52)	7	7	$P(^1P_1)_{ud}$		.067
MSTJ(101)	5	5	$P(^3P_0)_{ud}$		.056
MSTJ(107)	0	0	$P(^3P_1)_{ud}$		.067
PARJ(1)	0.10	0.10	$P(^3P_2)_{ud}$		.112
PARJ(2)	0.28	0.28	$P(^1S_0)_s$		.388
PARJ(3)	0.55	0.55	$P(^3S_1)_s$		.296
PARJ(4)	0.07	0.07	$P(^1P_1)_s$		.079
PARJ(5)	0.5	0.5	$P(^3P_0)_s$		.026
PARJ(11)	0.55		$P(^3P_1)_s$		.079
PARJ(12)	0.55		$P(^3P_2)_s$		.132
PARJ(13)	0.75		$P(^1S_0)_c$		.250
PARJ(14)	0.090		$P(^3S_1)_c$		.400
PARJ(15)	0.070		$P(^1P_1)_c$		.087
PARJ(16)	0.085		$P(^3P_0)_c$		.030
PARJ(17)	0.140		$P(^3P_1)_c$		.087
PARJ(19)	0.5	0.5	$P(^3P_2)_c$		.146
PARJ(21)	0.417	0.428	$P(^1S_0)_b$		.1625
PARJ(25)	0.7	0.7	$P(^3S_1)_b$		.4875
PARJ(26)	0.2	0.2	$P(^1P_1)_b$		.087
PARJ(41)	0.5	0.354	$P(^3P_0)_b$		.030
PARJ(42)	0.701	0.523	$P(^3P_1)_b$		.087
PARJ(54)	-0.0631	-0.0305	$P(^3P_2)_b$		.146
PARJ(55)	-0.00414	-0.00233			

Table 9: DELPHI parameter setting of JETSET 7.3 PS (December 1993 tuning used for modelling detector effects for 1993 data and September 1994 tuning used for modelling detector effects for 1994 data)

C Tables of Sensitivities

Parameter	JETSET 7.3 PS		JETSET 7.4 PS		ARIADNE 4.06		JETSET 7.4 ME	
	$\langle x_E \rangle_{D^*}$	$\langle x_E \rangle_B$	$\langle x_E \rangle_{D^*}$	$\langle x_E \rangle_B$	$\langle x_E \rangle_{D^*}$	$\langle x_E \rangle_B$	$\langle x_E \rangle_{D^*}$	$\langle x_E \rangle_B$
Λ_{QCD}	32	30	29	26	23	18	22	23
ϵ_C	9		9		10		9	
ϵ_B		8		8		9		10

Table 10: Sensitivity (x100) of $\langle x_E \rangle$ for D^* - and B-mesons

Property	a	b	σ_q	Λ_{QCD}	Q_0	γ_s	$\frac{P(qq)}{P(q)}$	u/d quarks		s quarks	
								$P(^1S_0)$	$P(^3S_1)$	$P(^1S_0)$	$P(^3S_1)$
S	1	2	5	10	1	4	2	1	1	0	0
A	2	6	12	20	1	9	12	1	2	4	1
P	1	2	2	8	1	2	1	1	1	0	1
$1 - T$	2	4	6	10	1	2	2	0	1	0	0
M	6	13	11	19	6	4	4	1	1	2	0
m	5	13	11	18	4	1	5	1	1	1	1
O	2	5	6	6	1	3	2	1	0	0	1
D_2^D	1	1	6	9	2	6	2	0	2	1	1
D_3^D	1	2	9	22	6	12	11	3	4	1	1
D_4^D	4	9	16	45	5	16	17	9	6	5	3
D_2^J	1	1	2	8	2	3	2	0	1	0	1
D_3^J	2	4	6	19	1	2	1	2	0	0	1
D_4^J	3	9	11	20	3	7	4	3	1	1	1
C	2	5	5	10	1	3	2	1	1	1	1
D	2	6	6	15	2	4	5	1	0	1	1
$M_h^2/E_{vis.}^2$	2	5	3	9	1	4	1	1	0	1	1
$M_l^2/E_{vis.}^2$	5	11	2	23	3	4	4	1	1	1	1
$M_d^2/E_{vis.}^2$	1	4	4	6	0	6	2	1	2	2	0
B_{max}	3	8	7	14	2	2	2	1	0	1	1
B_{min}	3	7	7	20	4	2	6	3	1	2	2
B_{sum}	5	10	8	15	4	5	4	1	1	1	0
$B_{diff.}$	1	4	5	5	1	5	3	1	2	2	1
p_t^{in}	5	8	9	5	3	3	3	2	2	2	1
p_t^{out}	4	8	27	11	2	2	2	4	2	2	0
y_T	2	5	6	8	2	1	2	1	1	1	0
x_p	3	6	7	6	4	1	1	2	1	1	0
$p_t^{out,vs.x_p}$	3	7	7	9	7	11	6	3	2	2	0
$p_t^{trans,vs.x_p}$	3	7	7	9	7	11	6	3	2	2	0
EEC	1	1	1	4	0	1	0	0	0	0	0
$AEEC$	1	6	16	9	13	32	17	6	2	7	3
ρ°	4	7	9	8	4	4	3	3	9	1	1
$K^0/K^\pm$	4	7	9	6	4	13	4	1	0	6	2
$K^{*0}/K^{*\pm}$	4	8	10	7	4	14	4	0	1	8	10
ϕ	5	8	7	9	4	35	9	3	3	5	23
p	5	12	13	5	6	3	22	3	1	1	3
Λ^0	5	11	12	12	4	17	21	2	2	2	2
$\Sigma^\pm(1193)$	4	15	18	11	8	9	21	4	4	5	5

Table 11: Sensitivities (x100) for JETSET 7.4 PS with standard decays. ARIADNE (with Q_0 in place of p_t^{QCD}) and JETSET 7.3 PS sensitivities are similar. The identified particle sensitivities have been calculated for the ξ_p distributions. The quoted values have significant statistical errors.

Property	<i>QCDLAM</i>	<i>RMASS</i> (13)	<i>CLMAX</i>	<i>CLPOW</i>	<i>CLSMR</i>	<i>PWT</i> (3)	<i>PWT</i> (7)	<i>DECWT</i>
<i>S</i>	11	3	2	3	1	1	0	1
<i>A</i>	25	3	7	5	2	2	1	3
<i>P</i>	9	2	2	2	0	0	1	1
$1 - T$	12	6	1	1	0	1	0	1
<i>M</i>	13	7	3	7	1	1	1	2
<i>m</i>	20	5	3	4	1	1	1	2
<i>O</i>	8	1	2	1	0	1	1	1
D_2^D	12	2	3	1	1	1	1	0
D_3^D	27	6	4	3	1	1	1	2
D_4^D	39	5	8	6	3	5	3	2
D_2^J	11	2	2	1	0	0	1	1
D_3^J	21	5	3	3	1	2	1	1
D_4^J	24	6	7	5	1	1	1	2
<i>C</i>	12	6	1	1	0	1	0	1
<i>D</i>	20	5	2	1	1	1	1	2
$M_b^2/E_{vis.}^2$	10	7	4	2	1	1	1	1
$M_l^2/E_{vis.}^2$	17	14	5	2	1	2	1	1
$M_d^2/E_{vis.}^2$	10	3	4	3	1	0	1	1
<i>B_{max}</i>	13	8	3	4	1	1	0	1
<i>B_{min}</i>	19	7	3	2	1	1	1	2
<i>B_{sum}</i>	12	7	1	4	1	0	1	2
<i>B_{diff.}</i>	8	1	2	1	0	1	1	1
p_t^{in}	6	4	5	11	3	1	0	1
p_t^{out}	15	5	14	7	2	1	0	2
y_T	6	4	7	10	2	1	1	1
x_p	2	3	4	9	3	1	1	0
$p_t^{out} vs. x_p$	1	3	3	50	5	2	2	5
$p_t^{trans.} vs. x_p$	1	3	3	50	5	2	2	5
<i>EEC</i>	5	1	1	0	0	0	0	0
<i>AEEC</i>	7	3	3	4	1	1	2	4
ρ^0	4	6	7	12	3	4	2	2
$K^0/K^\pm$	6	11	13	23	6	13	4	2
$K^{*0}/K^{*\pm}$	8	14	8	15	4	14	5	3
ϕ	15	24	10	21	6	33	10	3
<i>p</i>	5	7	48	20	7	2	3	15
Λ^0	9	13	65	27	8	18	8	11
$\Sigma^\pm(1193)$	10	18	46	16	7	13	7	7

Table 12: Sensitivities (x100) for HERWIG 5.8c with standard decays. The identified particle sensitivities have been calculated for the ξ_p distributions. The quoted values have significant statistical errors.

D Tables of Inclusive Charge Particle and Event Shape Distributions

The cross section has been determined from charged particles only and from charged and neutral particles. The tables show first the measurement from charged particles corrected to the charged final state, then corrected to the full final state. The third column contains the measurement from charged plus neutral particles corrected to the full final state. For inclusive charged particle distributions the relevant event axes have been evaluated for the corresponding final states. The cross section given in the table is followed by the statistical and systematic errors.

D.1 Tables of Inclusive Charged Particle Distributions

p_t^{in} with respect to the Thrust axis					$\frac{1}{N} \frac{dN}{dp_t^{in}}$				
Interval/GeV	charged particles corrected to charged particles				charged particles corrected to charged plus neutral particles				charged plus neutral particles corrected to charged plus neutral particles
0.000 - 0.100	44.667	± 0.036	± 1.653		44.740	± 0.036	± 1.685		46.663 ± 0.037 ± 1.758
0.100 - 0.200	39.331	± 0.033	± 1.062		39.475	± 0.033	± 1.078		39.823 ± 0.033 ± 1.092
0.200 - 0.300	29.429	± 0.028	± 0.603		29.491	± 0.028	± 0.606		29.351 ± 0.028 ± 0.608
0.300 - 0.400	21.326	± 0.024	± 0.353		21.338	± 0.024	± 0.352		21.034 ± 0.024 ± 0.350
0.400 - 0.500	15.547	± 0.020	± 0.225		15.463	± 0.020	± 0.222		15.156 ± 0.020 ± 0.219
0.500 - 0.600	11.463	± 0.017	± 0.156		11.437	± 0.017	± 0.154		11.149 ± 0.017 ± 0.150
0.600 - 0.700	8.668	± 0.015	± 0.116		8.589	± 0.015	± 0.115		8.348 ± 0.015 ± 0.111
0.700 - 0.800	6.664	± 0.013	± 0.090		6.608	± 0.013	± 0.091		6.430 ± 0.013 ± 0.087
0.800 - 1.000	4.6916	± 0.0078	± 0.0646		4.6518	± 0.0078	± 0.0661		4.5131 ± 0.0076 ± 0.0624
1.000 - 1.200	3.0694	± 0.0063	± 0.0430		3.0346	± 0.0063	± 0.0449		2.9522 ± 0.0062 ± 0.0420
1.200 - 1.400	2.0913	± 0.0052	± 0.0298		2.0847	± 0.0052	± 0.0322		2.0401 ± 0.0052 ± 0.0299
1.400 - 1.600	1.5030	± 0.0044	± 0.0218		1.4924	± 0.0044	± 0.0243		1.4597 ± 0.0044 ± 0.0222
1.600 - 1.800	1.0916	± 0.0038	± 0.0163		1.0961	± 0.0038	± 0.0189		1.0796 ± 0.0038 ± 0.0171
1.800 - 2.000	0.8306	± 0.0033	± 0.0130		0.8228	± 0.0033	± 0.0153		0.8155 ± 0.0033 ± 0.0136
2.000 - 2.500	0.5311	± 0.0017	± 0.0089		0.5315	± 0.0017	± 0.0108		0.5326 ± 0.0017 ± 0.0095
2.500 - 3.000	0.2946	± 0.0013	± 0.0055		0.2935	± 0.0013	± 0.0066		0.2988 ± 0.0013 ± 0.0057
3.000 - 3.500	0.17405	± 0.00097	± 0.00372		0.17393	± 0.00097	± 0.00441		0.18067 ± 0.00099 ± 0.00383
3.500 - 4.000	0.10909	± 0.00077	± 0.00276		0.10957	± 0.00077	± 0.00319		0.11471 ± 0.00079 ± 0.00273
4.000 - 5.000	0.05911	± 0.00041	± 0.00182		0.05897	± 0.00040	± 0.00200		0.06305 ± 0.00042 ± 0.00171
5.000 - 6.000	0.02669	± 0.00028	± 0.00101		0.02672	± 0.00027	± 0.00106		0.03040 ± 0.00029 ± 0.00095
6.000 - 7.000	0.01369	± 0.00021	± 0.00064		0.01299	± 0.00019	± 0.00060		0.01501 ± 0.00021 ± 0.00054
7.000 - 8.000	0.00675	± 0.00014	± 0.00039		0.00670	± 0.00013	± 0.00035		0.00858 ± 0.00016 ± 0.00035
8.000 - 10.000	0.00271	± 0.00006	± 0.00019		0.00278	± 0.00006	± 0.00016		0.00376 ± 0.00008 ± 0.00017
10.000 - 12.000	0.00095	± 0.00004	± 0.00008		0.00084	± 0.00003	± 0.00005		0.00123 ± 0.00004 ± 0.00006
12.000 - 14.000	0.00031	± 0.00002	± 0.00003		0.00030	± 0.00002	± 0.00002		0.00044 ± 0.00003 ± 0.00002

Table 13: Transverse momentum p_t^{in} with respect to the Thrust axis

p_t^{out} with respect to the Thrust axis					$\frac{1}{N} \frac{dN}{dp_t^{out}}$				
Interval/GeV	charged particles corrected to charged particles				charged particles corrected to charged plus neutral particles			charged plus neutral particles corrected to charged plus neutral particles	
0.000 - 0.100	67.512	± 0.043	± 1.956		67.673	± 0.044	± 1.818	66.160	± 0.043 ± 1.822
0.100 - 0.200	50.517	± 0.037	± 1.024		50.698	± 0.037	± 1.166	49.794	± 0.037 ± 1.149
0.200 - 0.300	33.618	± 0.030	± 0.593		33.803	± 0.030	± 0.692	33.544	± 0.030 ± 0.678
0.300 - 0.400	21.228	± 0.024	± 0.368		21.278	± 0.024	± 0.403	21.407	± 0.024 ± 0.397
0.400 - 0.500	13.160	± 0.019	± 0.235		13.171	± 0.019	± 0.240	13.466	± 0.019 ± 0.239
0.500 - 0.600	8.183	± 0.015	± 0.152		8.156	± 0.015	± 0.147	8.527	± 0.015 ± 0.150
0.600 - 0.700	5.178	± 0.012	± 0.101		5.126	± 0.011	± 0.094	5.448	± 0.012 ± 0.097
0.700 - 0.800	3.3455	± 0.0095	± 0.0684		3.3285	± 0.0093	± 0.0631	3.5845	± 0.0098 ± 0.0658
0.800 - 1.000	1.8640	± 0.0050	± 0.0407		1.8393	± 0.0049	± 0.0374	2.0309	± 0.0052 ± 0.0398
1.000 - 1.200	0.8873	± 0.0035	± 0.0210		0.8706	± 0.0034	± 0.0198	0.9959	± 0.0037 ± 0.0216
1.200 - 1.400	0.4600	± 0.0026	± 0.0117		0.4474	± 0.0024	± 0.0114	0.5288	± 0.0028 ± 0.0127
1.400 - 1.600	0.2500	± 0.0020	± 0.0068		0.2462	± 0.0018	± 0.0070	0.2987	± 0.0021 ± 0.0079
1.600 - 1.800	0.1429	± 0.0015	± 0.0042		0.1373	± 0.0014	± 0.0043	0.1755	± 0.0016 ± 0.0051
1.800 - 2.000	0.0841	± 0.0011	± 0.0026		0.0833	± 0.0011	± 0.0029	0.1086	± 0.0013 ± 0.0034
2.000 - 2.500	0.04069	± 0.00052	± 0.00140		0.03877	± 0.00047	± 0.00154	0.05266	± 0.00058 ± 0.00189
2.500 - 3.000	0.01468	± 0.00033	± 0.00057		0.01338	± 0.00028	± 0.00063	0.01885	± 0.00035 ± 0.00080
3.000 - 3.500	0.00501	± 0.00018	± 0.00022		0.00501	± 0.00017	± 0.00028	0.00814	± 0.00023 ± 0.00040

Table 14: Transverse momentum p_t^{out} with respect to the Thrust axis

p_t^{in} with respect to the Sphericity axis					$\frac{1}{N} \frac{dN}{dp_t^{in}}$				
Interval /GeV	charged particles corrected to charged particles				charged particles corrected to charged plus neutral particles			charged plus neutral particles corrected to charged plus neutral particles	
0.000 - 0.100	49.063	± 0.038	± 1.068		48.833	± 0.038	± 1.696	49.206	± 0.038 ± 1.672
0.100 - 0.200	38.338	± 0.032	± 0.819		38.301	± 0.033	± 1.265	38.461	± 0.033 ± 0.984
0.200 - 0.300	28.259	± 0.027	± 0.593		28.231	± 0.027	± 0.886	28.203	± 0.027 ± 0.571
0.300 - 0.400	20.492	± 0.023	± 0.422		20.554	± 0.023	± 0.613	20.391	± 0.023 ± 0.349
0.400 - 0.500	15.073	± 0.020	± 0.306		15.123	± 0.020	± 0.428	14.926	± 0.020 ± 0.233
0.500 - 0.600	11.274	± 0.017	± 0.225		11.312	± 0.017	± 0.304	11.133	± 0.017 ± 0.168
0.600 - 0.700	8.566	± 0.015	± 0.169		8.601	± 0.015	± 0.219	8.458	± 0.015 ± 0.129
0.700 - 0.800	6.627	± 0.013	± 0.129		6.659	± 0.013	± 0.161	6.548	± 0.013 ± 0.102
0.800 - 1.000	4.7065	± 0.0078	± 0.0898		4.7326	± 0.0079	± 0.1055	4.6706	± 0.0078 ± 0.0747
1.000 - 1.200	3.0724	± 0.0063	± 0.0574		3.0960	± 0.0064	± 0.0620	3.0684	± 0.0064 ± 0.0504
1.200 - 1.400	2.1120	± 0.0052	± 0.0388		2.1289	± 0.0053	± 0.0383	2.1299	± 0.0053 ± 0.0359
1.400 - 1.600	1.4873	± 0.0044	± 0.0270		1.5068	± 0.0045	± 0.0244	1.5201	± 0.0045 ± 0.0264
1.600 - 1.800	1.0882	± 0.0038	± 0.0196		1.0925	± 0.0038	± 0.0160	1.1143	± 0.0039 ± 0.0201
1.800 - 2.000	0.8101	± 0.0033	± 0.0146		0.8173	± 0.0033	± 0.0109	0.8398	± 0.0034 ± 0.0159
2.000 - 2.500	0.5156	± 0.0017	± 0.0094		0.5172	± 0.0017	± 0.0060	0.5334	± 0.0017 ± 0.0107
2.500 - 3.000	0.2799	± 0.0012	± 0.0053		0.2793	± 0.0012	± 0.0029	0.2968	± 0.0013 ± 0.0065
3.000 - 3.500	0.16206	± 0.00095	± 0.00323		0.16110	± 0.00093	± 0.00168	0.17343	± 0.00098 ± 0.00418
3.500 - 4.000	0.09848	± 0.00074	± 0.00212		0.09814	± 0.00073	± 0.00116	0.10741	± 0.00078 ± 0.00292
4.000 - 5.000	0.04987	± 0.00038	± 0.00124		0.04903	± 0.00036	± 0.00080	0.05615	± 0.00040 ± 0.00176
5.000 - 6.000	0.02113	± 0.00025	± 0.00065		0.02055	± 0.00024	± 0.00054	0.02473	± 0.00027 ± 0.00089
6.000 - 7.000	0.00933	± 0.00017	± 0.00036		0.00907	± 0.00016	± 0.00037	0.01157	± 0.00019 ± 0.00048
7.000 - 8.000	0.00401	± 0.00011	± 0.00019		0.00400	± 0.00011	± 0.00024	0.00561	± 0.00013 ± 0.00026
8.000 - 10.000	0.00132	± 0.00005	± 0.00008		0.00135	± 0.00004	± 0.00013	0.00204	± 0.00006 ± 0.00010
10.000 - 12.000	0.00026	± 0.00002	± 0.00002		0.00030	± 0.00002	± 0.00005	0.00049	± 0.00003 ± 0.00002
12.000 - 14.000	0.00007	± 0.00001	± 0.00001		0.00007	± 0.00001	± 0.00002	0.00012	± 0.00001 ± 0.00001

Table 15: Transverse momentum p_t^{in} with respect to the Sphericity axis

p_t^{out} with respect to the Sphericity axis					$\frac{1}{N} \frac{dN}{dp_t^{out}}$				
Interval/GeV	charged particles corrected to charged particles				charged particles corrected to charged plus neutral particles				charged plus neutral particles corrected to charged plus neutral particles
0.000 - 0.100	68.848	± 0.044	± 1.931		69.511	± 0.045	± 1.992		66.825 ± 0.043 ± 1.506
0.100 - 0.200	51.541	± 0.037	± 1.039		51.918	± 0.038	± 1.141		50.556 ± 0.037 ± 1.102
0.200 - 0.300	34.351	± 0.030	± 0.607		34.419	± 0.030	± 0.644		34.241 ± 0.030 ± 0.726
0.300 - 0.400	21.343	± 0.024	± 0.373		21.274	± 0.024	± 0.372		21.708 ± 0.024 ± 0.451
0.400 - 0.500	12.939	± 0.018	± 0.236		12.816	± 0.018	± 0.224		13.481 ± 0.019 ± 0.277
0.500 - 0.600	7.765	± 0.014	± 0.150		7.641	± 0.014	± 0.140		8.314 ± 0.015 ± 0.170
0.600 - 0.700	4.757	± 0.011	± 0.098		4.660	± 0.011	± 0.091		5.180 ± 0.012 ± 0.106
0.700 - 0.800	2.9543	± 0.0090	± 0.0651		2.8799	± 0.0085	± 0.0605		3.2986 ± 0.0094 ± 0.0679
0.800 - 1.000	1.5438	± 0.0047	± 0.0373		1.5015	± 0.0044	± 0.0354		1.7559 ± 0.0049 ± 0.0370
1.000 - 1.200	0.6885	± 0.0032	± 0.0186		0.6683	± 0.0030	± 0.0182		0.8187 ± 0.0034 ± 0.0181
1.200 - 1.400	0.3309	± 0.0023	± 0.0099		0.3218	± 0.0021	± 0.0100		0.4064 ± 0.0024 ± 0.0096
1.400 - 1.600	0.1730	± 0.0017	± 0.0057		0.1678	± 0.0015	± 0.0058		0.2175 ± 0.0018 ± 0.0055
1.600 - 1.800	0.0956	± 0.0013	± 0.0034		0.0909	± 0.0011	± 0.0035		0.1232 ± 0.0014 ± 0.0034
1.800 - 2.000	0.05183	± 0.00093	± 0.00199		0.05046	± 0.00085	± 0.00215		0.0712 ± 0.0011 ± 0.0022
2.000 - 2.500	0.02330	± 0.00041	± 0.00101		0.02209	± 0.00036	± 0.00109		0.03217 ± 0.00047 ± 0.00115
2.500 - 3.000	0.00729	± 0.00023	± 0.00037		0.00708	± 0.00021	± 0.00042		0.01112 ± 0.00029 ± 0.00050
3.000 - 3.500	0.00230	± 0.00013	± 0.00013		0.00219	± 0.00011	± 0.00015		0.00387 ± 0.00017 ± 0.00021

Table 16: Transverse momentum p_t^{out} with respect to the Sphericity axis

Rapidity y_T with respect to the Thrust axis					$\frac{1}{N} \frac{dN}{dy_T}$				
Interval/GeV	charged particles corrected to charged particles				charged particles corrected to charged plus neutral particles				charged plus neutral particles corrected to charged plus neutral particles
0.000 - 0.250	5.7889	± 0.0093	± 0.5509		5.7590	± 0.0093	± 0.5296		5.9517 ± 0.0095 ± 0.5628
0.250 - 0.500	6.3749	± 0.0094	± 0.4306		6.4046	± 0.0095	± 0.4411		6.4291 ± 0.0095 ± 0.4417
0.500 - 0.750	6.6630	± 0.0093	± 0.3197		6.6834	± 0.0094	± 0.3399		6.6831 ± 0.0094 ± 0.3319
0.750 - 1.000	6.7572	± 0.0089	± 0.2316		6.8119	± 0.0090	± 0.2522		6.7763 ± 0.0089 ± 0.2429
1.000 - 1.250	6.7638	± 0.0084	± 0.1677		6.8102	± 0.0085	± 0.1810		6.7650 ± 0.0085 ± 0.1755
1.250 - 1.500	6.7463	± 0.0082	± 0.1238		6.7853	± 0.0083	± 0.1284		6.7230 ± 0.0083 ± 0.1277
1.500 - 1.750	6.6488	± 0.0081	± 0.0933		6.6622	± 0.0082	± 0.0898		6.6085 ± 0.0081 ± 0.0939
1.750 - 2.000	6.5006	± 0.0080	± 0.0729		6.4845	± 0.0081	± 0.0648		6.4346 ± 0.0080 ± 0.0710
2.000 - 2.250	6.2303	± 0.0079	± 0.0623		6.1936	± 0.0079	± 0.0619		6.1697 ± 0.0078 ± 0.0617
2.250 - 2.500	5.8228	± 0.0076	± 0.0582		5.7642	± 0.0076	± 0.0576		5.7692 ± 0.0076 ± 0.0577
2.500 - 2.750	5.1930	± 0.0072	± 0.0519		5.1348	± 0.0072	± 0.0513		5.1450 ± 0.0072 ± 0.0514
2.750 - 3.000	4.3686	± 0.0067	± 0.0437		4.3142	± 0.0066	± 0.0431		4.3511 ± 0.0066 ± 0.0435
3.000 - 3.250	3.4544	± 0.0059	± 0.0345		3.4239	± 0.0059	± 0.0342		3.4481 ± 0.0059 ± 0.0345
3.250 - 3.500	2.5583	± 0.0051	± 0.0256		2.5473	± 0.0051	± 0.0255		2.5852 ± 0.0051 ± 0.0259
3.500 - 3.750	1.7687	± 0.0042	± 0.0177		1.7878	± 0.0043	± 0.0196		1.7999 ± 0.0043 ± 0.0180
3.750 - 4.000	1.1330	± 0.0034	± 0.0113		1.1585	± 0.0034	± 0.0150		1.1669 ± 0.0035 ± 0.0117
4.000 - 4.250	0.6770	± 0.0026	± 0.0068		0.7044	± 0.0027	± 0.0107		0.7054 ± 0.0027 ± 0.0071
4.250 - 4.500	0.3814	± 0.0019	± 0.0038		0.4054	± 0.0020	± 0.0071		0.3997 ± 0.0020 ± 0.0041
4.500 - 5.000	0.14699	± 0.00084	± 0.00147		0.16077	± 0.00091	± 0.00339		0.15673 ± 0.00089 ± 0.00177
5.000 - 5.500	0.03148	± 0.00038	± 0.00031		0.03466	± 0.00042	± 0.00091		0.03374 ± 0.00041 ± 0.00043
5.500 - 6.000	0.00486	± 0.00016	± 0.00005		0.00543	± 0.00017	± 0.00017		0.00502 ± 0.00016 ± 0.00007

Table 17: Rapidity y_T with respect to the Thrust axis

Rapidity y_S with respect to the Sphericity axis				$\frac{1}{N} \frac{dN}{dy_S}$			
Interval	charged particles corrected to charged particles			charged particles corrected to charged plus neutral particles			charged plus neutral particles corrected to charged plus neutral particles
0.000 - 0.250	6.5422	± 0.0096	± 0.5337	6.5778	± 0.0097	± 0.5082	6.5680 ± 0.0097 ± 0.5323
0.250 - 0.500	6.5743	± 0.0096	± 0.4132	6.6009	± 0.0096	± 0.4157	6.5901 ± 0.0096 ± 0.4246
0.500 - 0.750	6.5997	± 0.0092	± 0.3151	6.6327	± 0.0093	± 0.3348	6.6094 ± 0.0093 ± 0.3329
0.750 - 1.000	6.5896	± 0.0087	± 0.2357	6.6311	± 0.0088	± 0.2630	6.6152 ± 0.0088 ± 0.2554
1.000 - 1.250	6.5660	± 0.0083	± 0.1734	6.6019	± 0.0084	± 0.2011	6.5917 ± 0.0084 ± 0.1908
1.250 - 1.500	6.5623	± 0.0081	± 0.1266	6.5787	± 0.0082	± 0.1499	6.5817 ± 0.0082 ± 0.1393
1.500 - 1.750	6.4983	± 0.0080	± 0.0911	6.5086	± 0.0081	± 0.1079	6.5221 ± 0.0081 ± 0.0983
1.750 - 2.000	6.3665	± 0.0080	± 0.0656	6.3719	± 0.0080	± 0.0747	6.4097 ± 0.0080 ± 0.0673
2.000 - 2.250	6.1189	± 0.0078	± 0.0612	6.1073	± 0.0078	± 0.0611	6.1741 ± 0.0079 ± 0.0617
2.250 - 2.500	5.6636	± 0.0076	± 0.0566	5.6636	± 0.0076	± 0.0566	5.7542 ± 0.0076 ± 0.0575
2.500 - 2.750	5.0282	± 0.0071	± 0.0503	5.0161	± 0.0071	± 0.0502	5.1066 ± 0.0072 ± 0.0511
2.750 - 3.000	4.2102	± 0.0066	± 0.0421	4.1948	± 0.0065	± 0.0419	4.2721 ± 0.0066 ± 0.0427
3.000 - 3.250	3.3399	± 0.0058	± 0.0334	3.3206	± 0.0058	± 0.0332	3.3718 ± 0.0059 ± 0.0337
3.250 - 3.500	2.5260	± 0.0051	± 0.0253	2.4999	± 0.0050	± 0.0250	2.5185 ± 0.0051 ± 0.0252
3.500 - 3.750	1.7861	± 0.0042	± 0.0179	1.7772	± 0.0043	± 0.0178	1.7588 ± 0.0042 ± 0.0176
3.750 - 4.000	1.2194	± 0.0035	± 0.0142	1.2049	± 0.0035	± 0.0155	1.1589 ± 0.0034 ± 0.0130
4.000 - 4.250	0.8020	± 0.0028	± 0.0110	0.7860	± 0.0028	± 0.0133	0.7327 ± 0.0027 ± 0.0105
4.250 - 4.500	0.4995	± 0.0022	± 0.0080	0.4933	± 0.0022	± 0.0106	0.4402 ± 0.0021 ± 0.0078
4.500 - 5.000	0.2475	± 0.0011	± 0.0049	0.2406	± 0.0011	± 0.0071	0.1952 ± 0.0010 ± 0.0046
5.000 - 5.500	0.08078	± 0.00060	± 0.00203	0.07713	± 0.00064	± 0.00326	0.05574 ± 0.00050 ± 0.00180
5.500 - 6.000	0.02382	± 0.00032	± 0.00073	0.02004	± 0.00033	± 0.00113	0.01306 ± 0.00024 ± 0.00055

Table 18: Rapidity y_S with respect to the Sphericity axis

Scaled momentum x_p	$\frac{1}{N} \frac{dN}{dx_p}$		
Interval	charged particles		
	corrected to charged particles		
0.000 - 0.010	389.18	± 0.37	± 28.68
0.010 - 0.020	407.47	± 0.32	± 10.44
0.020 - 0.030	266.10	± 0.26	± 3.67
0.030 - 0.040	185.45	± 0.22	± 1.85
0.040 - 0.050	137.73	± 0.19	± 1.38
0.050 - 0.060	106.07	± 0.16	± 1.08
0.060 - 0.070	84.57	± 0.15	± 0.95
0.070 - 0.080	68.87	± 0.13	± 0.85
0.080 - 0.090	57.09	± 0.12	± 0.78
0.090 - 0.100	47.79	± 0.11	± 0.71
0.100 - 0.120	37.756	± 0.071	± 0.619
0.120 - 0.140	28.006	± 0.061	± 0.501
0.140 - 0.160	21.330	± 0.054	± 0.409
0.160 - 0.180	16.660	± 0.048	± 0.337
0.180 - 0.200	13.043	± 0.042	± 0.273
0.200 - 0.250	9.019	± 0.022	± 0.195
0.250 - 0.300	5.464	± 0.018	± 0.124
0.300 - 0.400	2.8036	± 0.0088	± 0.0683
0.400 - 0.500	1.1938	± 0.0057	± 0.0327
0.500 - 0.600	0.5322	± 0.0040	± 0.0169
0.600 - 0.700	0.2287	± 0.0027	± 0.0084
0.700 - 0.800	0.0936	± 0.0016	± 0.0039
0.800 - 1.000	0.0201	± 0.0005	± 0.0023

Table 19: Scaled momentum x_p

$\xi_p = \ln(\frac{1}{x_p})$	$\frac{1}{N} \frac{dN}{d\xi_p}$		
Interval	charged particles		
	corrected to charged particles		
0.000 - 0.200	0.0151	± 0.0004	± 0.0022
0.200 - 0.400	0.0774	± 0.0011	± 0.0051
0.400 - 0.600	0.1987	± 0.0018	± 0.0083
0.600 - 0.800	0.3970	± 0.0023	± 0.0116
0.800 - 1.000	0.6805	± 0.0031	± 0.0161
1.000 - 1.200	1.0520	± 0.0038	± 0.0226
1.200 - 1.400	1.5128	± 0.0046	± 0.0311
1.400 - 1.600	2.0270	± 0.0053	± 0.0398
1.600 - 1.800	2.5997	± 0.0060	± 0.0481
1.800 - 2.000	3.1993	± 0.0066	± 0.0553
2.000 - 2.200	3.8069	± 0.0071	± 0.0610
2.200 - 2.400	4.3851	± 0.0076	± 0.0651
2.400 - 2.600	4.9235	± 0.0080	± 0.0680
2.600 - 2.800	5.4160	± 0.0083	± 0.0702
2.800 - 3.000	5.8079	± 0.0086	± 0.0716
3.000 - 3.200	6.1608	± 0.0088	± 0.0734
3.200 - 3.400	6.4026	± 0.0090	± 0.0753
3.400 - 3.600	6.5371	± 0.0091	± 0.0779
3.600 - 3.800	6.5822	± 0.0091	± 0.0819
3.800 - 4.000	6.4738	± 0.0090	± 0.0869
4.000 - 4.200	6.2383	± 0.0089	± 0.0933
4.200 - 4.400	5.8490	± 0.0087	± 0.1003
4.400 - 4.600	5.3231	± 0.0084	± 0.1068
4.600 - 4.800	4.6603	± 0.0079	± 0.1098
4.800 - 5.000	3.9029	± 0.0073	± 0.1064
5.000 - 5.200	3.1150	± 0.0066	± 0.0952
5.200 - 5.400	2.3781	± 0.0058	± 0.0777

Table 20: $\xi_p = \ln(\frac{1}{x_p})$

x_p	$\langle p_t^{out} \rangle / GeV$		
Interval	charged particles		
	corrected to charged particles		
0.000 - 0.010	0.11124	± 0.00012	± 0.00097
0.010 - 0.020	0.19506	± 0.00014	± 0.00113
0.020 - 0.030	0.23694	± 0.00019	± 0.00118
0.030 - 0.040	0.25986	± 0.00023	± 0.00130
0.040 - 0.050	0.27485	± 0.00028	± 0.00137
0.050 - 0.060	0.28678	± 0.00032	± 0.00143
0.060 - 0.070	0.29812	± 0.00037	± 0.00149
0.070 - 0.080	0.30653	± 0.00041	± 0.00153
0.080 - 0.090	0.31324	± 0.00046	± 0.00157
0.090 - 0.100	0.31969	± 0.00051	± 0.00160
0.100 - 0.120	0.32899	± 0.00041	± 0.00164
0.120 - 0.140	0.34140	± 0.00049	± 0.00171
0.140 - 0.160	0.35107	± 0.00058	± 0.00176
0.160 - 0.180	0.35833	± 0.00066	± 0.00179
0.180 - 0.200	0.37165	± 0.00076	± 0.00186
0.200 - 0.250	0.38189	± 0.00059	± 0.00191
0.250 - 0.300	0.40047	± 0.00078	± 0.00200
0.300 - 0.400	0.41720	± 0.00079	± 0.00223
0.400 - 0.500	0.4348	± 0.0012	± 0.0031
0.500 - 0.600	0.4436	± 0.0019	± 0.0044
0.600 - 0.700	0.4412	± 0.0031	± 0.0062
0.700 - 0.800	0.4525	± 0.0046	± 0.0093
0.800 - 1.000	0.3778	± 0.0051	± 0.0112

Table 21: $\langle p_t^{out} \rangle$ vs. x_p

x_p	$\langle p_t \rangle / GeV$		
Interval	charged particles		
	corrected to charged particles		
0.000 - 0.010	0.18791	± 0.00016	± 0.00171
0.010 - 0.020	0.36450	± 0.00019	± 0.00214
0.020 - 0.030	0.48985	± 0.00027	± 0.00245
0.030 - 0.040	0.57682	± 0.00035	± 0.00288
0.040 - 0.050	0.64402	± 0.00043	± 0.00322
0.050 - 0.060	0.69974	± 0.00051	± 0.00350
0.060 - 0.070	0.75195	± 0.00059	± 0.00376
0.070 - 0.080	0.79370	± 0.00067	± 0.00397
0.080 - 0.090	0.83170	± 0.00075	± 0.00416
0.090 - 0.100	0.85958	± 0.00083	± 0.00430
0.100 - 0.120	0.90697	± 0.00069	± 0.00453
0.120 - 0.140	0.96796	± 0.00083	± 0.00484
0.140 - 0.160	1.01237	± 0.00098	± 0.00506
0.160 - 0.180	1.0555	± 0.0011	± 0.0053
0.180 - 0.200	1.0903	± 0.0013	± 0.0055
0.200 - 0.250	1.1594	± 0.0010	± 0.0061
0.250 - 0.300	1.2363	± 0.0014	± 0.0076
0.300 - 0.400	1.3226	± 0.0014	± 0.0100
0.400 - 0.500	1.4043	± 0.0022	± 0.0135
0.500 - 0.600	1.4327	± 0.0035	± 0.0183
0.600 - 0.700	1.4210	± 0.0056	± 0.0246
0.700 - 0.800	1.4134	± 0.0084	± 0.0336
0.800 - 1.000	1.226	± 0.011	± 0.040

Table 22: $\langle p_t \rangle$ vs. x_p

Energy Energy Correlation EEC	
Interval $\cos \chi$	charged particles corrected to charged particles
-1.000 - -0.960	3.0850 $\pm$ 0.0085 $\pm$ 0.0154
-0.960 - -0.920	0.7097 $\pm$ 0.0040 $\pm$ 0.0035
-0.920 - -0.880	0.3794 $\pm$ 0.0029 $\pm$ 0.0019
-0.880 - -0.840	0.2517 $\pm$ 0.0024 $\pm$ 0.0013
-0.840 - -0.800	0.18476 $\pm$ 0.00202 $\pm$ 0.00092
-0.800 - -0.760	0.14645 $\pm$ 0.00179 $\pm$ 0.00073
-0.760 - -0.720	0.11995 $\pm$ 0.00161 $\pm$ 0.00060
-0.720 - -0.680	0.10155 $\pm$ 0.00149 $\pm$ 0.00051
-0.680 - -0.640	0.08866 $\pm$ 0.00140 $\pm$ 0.00044
-0.640 - -0.600	0.07845 $\pm$ 0.00132 $\pm$ 0.00039
-0.600 - -0.560	0.07092 $\pm$ 0.00126 $\pm$ 0.00036
-0.560 - -0.520	0.06498 $\pm$ 0.00122 $\pm$ 0.00037
-0.520 - -0.480	0.05975 $\pm$ 0.00117 $\pm$ 0.00037
-0.480 - -0.440	0.05559 $\pm$ 0.00113 $\pm$ 0.00038
-0.440 - -0.400	0.05211 $\pm$ 0.00109 $\pm$ 0.00038
-0.400 - -0.360	0.04894 $\pm$ 0.00106 $\pm$ 0.00038
-0.360 - -0.320	0.04681 $\pm$ 0.00104 $\pm$ 0.00038
-0.320 - -0.280	0.04499 $\pm$ 0.00102 $\pm$ 0.00039
-0.280 - -0.240	0.04383 $\pm$ 0.00102 $\pm$ 0.00039
-0.240 - -0.200	0.04212 $\pm$ 0.00100 $\pm$ 0.00039
-0.200 - -0.160	0.04078 $\pm$ 0.00098 $\pm$ 0.00039
-0.160 - -0.120	0.03970 $\pm$ 0.00096 $\pm$ 0.00039
-0.120 - -0.080	0.03900 $\pm$ 0.00096 $\pm$ 0.00039
-0.080 - -0.040	0.03882 $\pm$ 0.00096 $\pm$ 0.00040
-0.040 - 0.000	0.03849 $\pm$ 0.00096 $\pm$ 0.00040
0.000 - 0.040	0.03825 $\pm$ 0.00096 $\pm$ 0.00040
0.040 - 0.080	0.03803 $\pm$ 0.00095 $\pm$ 0.00040
0.080 - 0.120	0.03816 $\pm$ 0.00096 $\pm$ 0.00040
0.120 - 0.160	0.03812 $\pm$ 0.00095 $\pm$ 0.00040
0.160 - 0.200	0.03874 $\pm$ 0.00096 $\pm$ 0.00041
0.200 - 0.240	0.03943 $\pm$ 0.00097 $\pm$ 0.00041
0.240 - 0.280	0.04023 $\pm$ 0.00098 $\pm$ 0.00041
0.280 - 0.320	0.04090 $\pm$ 0.00099 $\pm$ 0.00041
0.320 - 0.360	0.04233 $\pm$ 0.00101 $\pm$ 0.00041
0.360 - 0.400	0.04361 $\pm$ 0.00102 $\pm$ 0.00041
0.400 - 0.440	0.04521 $\pm$ 0.00103 $\pm$ 0.00040
0.440 - 0.480	0.04759 $\pm$ 0.00106 $\pm$ 0.00040
0.480 - 0.520	0.05050 $\pm$ 0.00109 $\pm$ 0.00040
0.520 - 0.560	0.05331 $\pm$ 0.00111 $\pm$ 0.00039
0.560 - 0.600	0.05702 $\pm$ 0.00114 $\pm$ 0.00038
0.600 - 0.640	0.06238 $\pm$ 0.00119 $\pm$ 0.00038
0.640 - 0.680	0.06943 $\pm$ 0.00126 $\pm$ 0.00038
0.680 - 0.720	0.07782 $\pm$ 0.00133 $\pm$ 0.00039
0.720 - 0.760	0.08949 $\pm$ 0.00142 $\pm$ 0.00045
0.760 - 0.800	0.10526 $\pm$ 0.00153 $\pm$ 0.00053
0.800 - 0.840	0.12980 $\pm$ 0.00170 $\pm$ 0.00065
0.840 - 0.880	0.17143 $\pm$ 0.00196 $\pm$ 0.00086
0.880 - 0.920	0.2539 $\pm$ 0.0024 $\pm$ 0.0013
0.920 - 0.960	0.4860 $\pm$ 0.0033 $\pm$ 0.0024
0.960 - 1.000	2.8965 $\pm$ 0.0083 $\pm$ 0.0166

Table 23: Energy Energy Correlation EEC

Asymmetry of the Energy Energy Correlation $AEEC$	
Interval $\cos \chi$	charged particles corrected to charged particles
-1.000 - -0.960	0.18959 $\pm$ 0.00051 $\pm$ 0.00924
-0.960 - -0.920	0.22480 $\pm$ 0.00150 $\pm$ 0.00220
-0.920 - -0.880	0.12560 $\pm$ 0.00120 $\pm$ 0.00130
-0.880 - -0.840	0.08023 $\pm$ 0.00092 $\pm$ 0.00080
-0.840 - -0.800	0.05496 $\pm$ 0.00073 $\pm$ 0.00055
-0.800 - -0.760	0.04122 $\pm$ 0.00062 $\pm$ 0.00049
-0.760 - -0.720	0.03054 $\pm$ 0.00050 $\pm$ 0.00046
-0.720 - -0.680	0.02383 $\pm$ 0.00042 $\pm$ 0.00044
-0.680 - -0.640	0.01929 $\pm$ 0.00036 $\pm$ 0.00043
-0.640 - -0.600	0.01613 $\pm$ 0.00032 $\pm$ 0.00042
-0.600 - -0.560	0.01391 $\pm$ 0.00029 $\pm$ 0.00041
-0.560 - -0.520	0.01182 $\pm$ 0.00026 $\pm$ 0.00079
-0.520 - -0.480	0.00938 $\pm$ 0.00021 $\pm$ 0.00036
-0.480 - -0.440	0.00803 $\pm$ 0.00019 $\pm$ 0.00035
-0.440 - -0.400	0.00696 $\pm$ 0.00017 $\pm$ 0.00033
-0.400 - -0.360	0.00546 $\pm$ 0.00014 $\pm$ 0.00029
-0.360 - -0.320	0.00459 $\pm$ 0.00012 $\pm$ 0.00027
-0.320 - -0.280	0.00408 $\pm$ 0.00011 $\pm$ 0.00026
-0.280 - -0.240	0.00360 $\pm$ 0.00010 $\pm$ 0.00025
-0.240 - -0.200	0.00268 $\pm$ 0.00008 $\pm$ 0.00020
-0.200 - -0.160	0.00212 $\pm$ 0.00006 $\pm$ 0.00017
-0.160 - -0.120	0.00148 $\pm$ 0.00005 $\pm$ 0.00013
-0.120 - -0.080	0.00104 $\pm$ 0.00003 $\pm$ 0.00010
-0.080 - -0.040	0.00069 $\pm$ 0.00003 $\pm$ 0.00010
-0.040 - 0.000	0.00013 $\pm$ 0.00002 $\pm$ 0.00005

Table 24: Asymmetry of the Energy Energy Correlation $AEEC$

D.2 Tables of Event Shape Distributions

Sphericity	$\frac{1}{N} \frac{dN}{dS}$								
Interval	charged particles corrected to charged particles			charged particles corrected to charged plus neutral particles			charged plus neutral particles corrected to charged plus neutral particles		
0.000 - 0.010	17.598	± 0.068	± 0.176	17.956	± 0.072	± 0.458	16.198	± 0.067	± 0.208
0.010 - 0.020	18.570	± 0.068	± 0.186	19.635	± 0.071	± 0.267	20.008	± 0.072	± 0.246
0.020 - 0.030	12.129	± 0.053	± 0.121	12.249	± 0.054	± 0.122	12.896	± 0.056	± 0.153
0.030 - 0.040	8.225	± 0.044	± 0.082	7.985	± 0.042	± 0.080	8.237	± 0.043	± 0.094
0.040 - 0.050	5.825	± 0.036	± 0.058	5.686	± 0.036	± 0.057	5.885	± 0.037	± 0.065
0.050 - 0.060	4.600	± 0.033	± 0.046	4.345	± 0.031	± 0.046	4.458	± 0.032	± 0.048
0.060 - 0.080	3.274	± 0.019	± 0.033	3.155	± 0.019	± 0.040	3.272	± 0.019	± 0.034
0.080 - 0.100	2.309	± 0.016	± 0.023	2.256	± 0.016	± 0.032	2.290	± 0.016	± 0.023
0.100 - 0.120	1.692	± 0.014	± 0.017	1.666	± 0.014	± 0.026	1.699	± 0.014	± 0.017
0.120 - 0.160	1.2001	± 0.0082	± 0.0132	1.1914	± 0.0082	± 0.0203	1.2018	± 0.0082	± 0.0120
0.160 - 0.200	0.8053	± 0.0067	± 0.0104	0.7994	± 0.0067	± 0.0145	0.7988	± 0.0067	± 0.0080
0.200 - 0.250	0.5688	± 0.0050	± 0.0084	0.5758	± 0.0051	± 0.0107	0.5610	± 0.0050	± 0.0063
0.250 - 0.300	0.4031	± 0.0043	± 0.0065	0.3972	± 0.0042	± 0.0071	0.3926	± 0.0042	± 0.0051
0.300 - 0.350	0.2931	± 0.0036	± 0.0049	0.2908	± 0.0036	± 0.0044	0.2810	± 0.0035	± 0.0043
0.350 - 0.400	0.2206	± 0.0031	± 0.0035	0.2244	± 0.0032	± 0.0023	0.2099	± 0.0030	± 0.0037
0.400 - 0.500	0.1528	± 0.0018	± 0.0025	0.1500	± 0.0018	± 0.0015	0.1441	± 0.0018	± 0.0032
0.500 - 0.600	0.0886	± 0.0014	± 0.0021	0.0900	± 0.0014	± 0.0026	0.0842	± 0.0013	± 0.0023
0.600 - 0.700	0.04319	± 0.00094	± 0.00209	0.04351	± 0.00095	± 0.00394	0.04160±	0.00092	± 0.00129
0.700 - 0.850	0.00778	± 0.00032	± 0.00081	0.00809	± 0.00034	± 0.00184	0.00758±	0.00032	± 0.00024

Table 25: Sphericity S

Aplanarity	$\frac{1}{N} \frac{dN}{dA}$								
Interval	charged particles corrected to charged particles			charged particles corrected to charged plus neutral particles			charged plus neutral particles corrected to charged plus neutral particles		
0.000 - 0.005	81.41	± 0.20	± 0.81	81.83	± 0.21	± 1.03	75.10	± 0.19	± 0.75
0.005 - 0.010	49.12	± 0.15	± 0.49	51.34	± 0.16	± 0.62	55.31	± 0.17	± 0.55
0.010 - 0.015	24.05	± 0.11	± 0.24	24.15	± 0.10	± 0.33	26.03	± 0.11	± 0.28
0.015 - 0.020	13.668	± 0.079	± 0.137	13.242	± 0.076	± 0.224	13.927	± 0.079	± 0.176
0.020 - 0.030	6.893	± 0.039	± 0.088	6.477	± 0.037	± 0.142	6.768	± 0.038	± 0.098
0.030 - 0.040	3.270	± 0.027	± 0.057	3.013	± 0.025	± 0.087	3.014	± 0.025	± 0.056
0.040 - 0.060	1.443	± 0.013	± 0.032	1.295	± 0.012	± 0.048	1.281	± 0.012	± 0.035
0.060 - 0.080	0.5841	± 0.0080	± 0.0153	0.5404	± 0.0077	± 0.0248	0.5181	± 0.0075	± 0.0188
0.080 - 0.100	0.2825	± 0.0056	± 0.0085	0.2687	± 0.0055	± 0.0150	0.2619	± 0.0054	± 0.0118
0.100 - 0.120	0.1515	± 0.0041	± 0.0051	0.1465	± 0.0041	± 0.0098	0.1461	± 0.0041	± 0.0079
0.120 - 0.140	0.0831	± 0.0030	± 0.0031	0.0797	± 0.0030	± 0.0061	0.0758	± 0.0029	± 0.0043
0.140 - 0.160	0.0535	± 0.0026	± 0.0023	0.0511	± 0.0025	± 0.0040	0.0467	± 0.0023	± 0.0027
0.160 - 0.200	0.0262	± 0.0012	± 0.0014	0.0239	± 0.0012	± 0.0017	0.0234	± 0.0011	± 0.0014
0.200 - 0.250	0.00969	± 0.00067	± 0.00062	0.01040	± 0.00071	± 0.00058	0.00884±	0.00061	± 0.00052
0.250 - 0.300	0.00320	± 0.00037	± 0.00075	0.00263	± 0.00035	± 0.00009	0.00310±	0.00040	± 0.00018

Table 26: Aplanarity A

Planarity				$\frac{1}{N} \frac{dN}{dP}$			
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles
	charged particles			charged plus neutral particles			corrected to charged plus neutral particles
0.000 - 0.005	64.81	± 0.18	± 0.65	68.07	± 0.19	± 0.82	68.69 ± 0.19 ± 0.74
0.005 - 0.010	32.31	± 0.12	± 0.32	31.44	± 0.12	± 0.39	31.66 ± 0.12 ± 0.35
0.010 - 0.015	18.177	± 0.092	± 0.182	17.196	± 0.088	± 0.215	17.091 ± 0.088 ± 0.188
0.015 - 0.020	12.001	± 0.075	± 0.120	11.362	± 0.072	± 0.145	11.370 ± 0.072 ± 0.127
0.020 - 0.025	8.607	± 0.062	± 0.086	8.373	± 0.062	± 0.109	8.417 ± 0.062 ± 0.095
0.025 - 0.030	6.792	± 0.056	± 0.068	6.542	± 0.055	± 0.087	6.578 ± 0.055 ± 0.075
0.030 - 0.035	5.542	± 0.051	± 0.055	5.414	± 0.050	± 0.074	5.479 ± 0.050 ± 0.063
0.035 - 0.040	4.519	± 0.045	± 0.045	4.461	± 0.045	± 0.062	4.493 ± 0.045 ± 0.052
0.040 - 0.050	3.640	± 0.029	± 0.036	3.585	± 0.029	± 0.051	3.610 ± 0.029 ± 0.042
0.050 - 0.060	2.757	± 0.025	± 0.028	2.747	± 0.025	± 0.041	2.749 ± 0.025 ± 0.033
0.060 - 0.080	1.986	± 0.015	± 0.022	1.986	± 0.015	± 0.031	1.987 ± 0.015 ± 0.024
0.080 - 0.100	1.362	± 0.012	± 0.017	1.394	± 0.013	± 0.023	1.362 ± 0.012 ± 0.017
0.100 - 0.120	1.008	± 0.011	± 0.014	1.026	± 0.011	± 0.018	1.008 ± 0.011 ± 0.013
0.120 - 0.160	0.6777	± 0.0061	± 0.0107	0.6898	± 0.0062	± 0.0133	0.6676 ± 0.0061 ± 0.0093
0.160 - 0.200	0.4477	± 0.0050	± 0.0083	0.4502	± 0.0051	± 0.0096	0.4248 ± 0.0048 ± 0.0063
0.200 - 0.250	0.2755	± 0.0034	± 0.0060	0.2807	± 0.0035	± 0.0067	0.2692 ± 0.0034 ± 0.0042
0.250 - 0.300	0.1801	± 0.0028	± 0.0045	0.1833	± 0.0029	± 0.0048	0.1742 ± 0.0028 ± 0.0029
0.300 - 0.350	0.1114	± 0.0022	± 0.0032	0.1121	± 0.0022	± 0.0033	0.1042 ± 0.0021 ± 0.0019
0.350 - 0.400	0.0575	± 0.0015	± 0.0018	0.0598	± 0.0016	± 0.0019	0.0566 ± 0.0015 ± 0.0011
0.400 - 0.500	0.0137	± 0.0005	± 0.0005	0.0144	± 0.0006	± 0.0005	0.0145 ± 0.0006 ± 0.0003

Table 27: Planarity P

1-Thrust				$\frac{1}{N} \frac{dN}{d(1-T)}$			
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles
	charged particles			charged plus neutral particles			corrected to charged plus neutral particles
0.000 - 0.010	3.386	± 0.028	± 0.132	1.366	± 0.025	± 0.125	1.030 ± 0.019 ± 0.076
0.010 - 0.020	13.901	± 0.057	± 0.249	10.185	± 0.048	± 0.446	10.951 ± 0.051 ± 0.527
0.020 - 0.030	15.733	± 0.061	± 0.157	16.184	± 0.062	± 0.288	17.645 ± 0.066 ± 0.547
0.030 - 0.040	12.535	± 0.055	± 0.125	13.572	± 0.059	± 0.173	14.192 ± 0.061 ± 0.292
0.040 - 0.050	9.494	± 0.048	± 0.095	10.004	± 0.050	± 0.134	10.009 ± 0.050 ± 0.152
0.050 - 0.060	7.248	± 0.042	± 0.072	7.581	± 0.044	± 0.108	7.572 ± 0.044 ± 0.101
0.060 - 0.070	5.665	± 0.037	± 0.057	5.933	± 0.039	± 0.090	5.760 ± 0.038 ± 0.076
0.070 - 0.080	4.544	± 0.033	± 0.045	4.723	± 0.035	± 0.072	4.619 ± 0.034 ± 0.062
0.080 - 0.090	3.722	± 0.030	± 0.040	3.929	± 0.032	± 0.056	3.792 ± 0.031 ± 0.051
0.090 - 0.100	3.085	± 0.027	± 0.038	3.265	± 0.029	± 0.043	3.176 ± 0.028 ± 0.042
0.100 - 0.120	2.413	± 0.017	± 0.027	2.578	± 0.018	± 0.035	2.456 ± 0.018 ± 0.032
0.120 - 0.140	1.800	± 0.015	± 0.018	1.916	± 0.016	± 0.029	1.825 ± 0.015 ± 0.022
0.140 - 0.160	1.387	± 0.013	± 0.014	1.477	± 0.014	± 0.025	1.401 ± 0.013 ± 0.016
0.160 - 0.180	1.074	± 0.011	± 0.011	1.142	± 0.012	± 0.020	1.074 ± 0.011 ± 0.011
0.180 - 0.200	0.8362	± 0.0099	± 0.0096	0.889	± 0.011	± 0.016	0.8262 ± 0.0100 ± 0.0083
0.200 - 0.250	0.5683	± 0.0051	± 0.0101	0.6005	± 0.0055	± 0.0123	0.5525 ± 0.0051 ± 0.0065
0.250 - 0.300	0.3138	± 0.0038	± 0.0077	0.3295	± 0.0040	± 0.0083	0.3030 ± 0.0038 ± 0.0058
0.300 - 0.350	0.1286	± 0.0024	± 0.0045	0.1395	± 0.0026	± 0.0062	0.1312 ± 0.0025 ± 0.0044
0.350 - 0.400	0.0180	± 0.0009	± 0.0011	0.0232	± 0.0012	± 0.0026	0.0238 ± 0.0012 ± 0.0014
0.400 - 0.500	0.0004	± 0.0001	± 0.0001	0.0005	± 0.0002	± 0.0002	0.0007 ± 0.0002 ± 0.0001

Table 28: 1-Thrust $(1 - T)$

Major	$\frac{1}{N} \frac{dN}{dM}$					
Interval	charged particles corrected to charged particles		charged particles corrected to charged plus neutral particles		charged plus neutral particles corrected to charged plus neutral particles	
0.000 - 0.020	0.00207	$\pm 0.00087 \pm 0.00313$	0.0022	$\pm 0.0023 \pm 0.0016$	0.00040	$\pm 0.00090 \pm 0.00005$
0.020 - 0.040	0.2165	$\pm 0.0053 \pm 0.0114$	0.1274	$\pm 0.0058 \pm 0.0094$	0.0590	$\pm 0.0030 \pm 0.0058$
0.040 - 0.050	1.300	$\pm 0.018 \pm 0.013$	0.895	$\pm 0.017 \pm 0.024$	0.642	$\pm 0.013 \pm 0.028$
0.050 - 0.060	2.734	$\pm 0.027 \pm 0.027$	2.337	$\pm 0.026 \pm 0.059$	2.178	$\pm 0.024 \pm 0.086$
0.060 - 0.070	4.245	$\pm 0.033 \pm 0.042$	4.132	$\pm 0.033 \pm 0.098$	4.303	$\pm 0.034 \pm 0.155$
0.070 - 0.080	5.280	$\pm 0.037 \pm 0.053$	5.469	$\pm 0.038 \pm 0.122$	5.849	$\pm 0.039 \pm 0.192$
0.080 - 0.100	6.130	$\pm 0.027 \pm 0.061$	6.370	$\pm 0.028 \pm 0.128$	6.889	$\pm 0.030 \pm 0.194$
0.100 - 0.120	5.861	$\pm 0.027 \pm 0.059$	5.870	$\pm 0.026 \pm 0.103$	6.342	$\pm 0.028 \pm 0.143$
0.120 - 0.140	4.881	$\pm 0.024 \pm 0.049$	4.726	$\pm 0.023 \pm 0.072$	4.890	$\pm 0.024 \pm 0.085$
0.140 - 0.160	3.966	$\pm 0.022 \pm 0.040$	3.876	$\pm 0.021 \pm 0.052$	3.900	$\pm 0.021 \pm 0.050$
0.160 - 0.200	3.024	$\pm 0.013 \pm 0.030$	3.002	$\pm 0.013 \pm 0.033$	2.960	$\pm 0.013 \pm 0.030$
0.200 - 0.240	2.166	$\pm 0.011 \pm 0.023$	2.169	$\pm 0.011 \pm 0.022$	2.124	$\pm 0.011 \pm 0.021$
0.240 - 0.280	1.5714	$\pm 0.0095 \pm 0.0177$	1.6042	$\pm 0.0097 \pm 0.0160$	1.5562	$\pm 0.0095 \pm 0.0156$
0.280 - 0.320	1.1782	$\pm 0.0082 \pm 0.0146$	1.2190	$\pm 0.0085 \pm 0.0122$	1.1807	$\pm 0.0083 \pm 0.0118$
0.320 - 0.360	0.8962	$\pm 0.0071 \pm 0.0125$	0.9228	$\pm 0.0074 \pm 0.0111$	0.8693	$\pm 0.0071 \pm 0.0087$
0.360 - 0.400	0.6771	$\pm 0.0061 \pm 0.0106$	0.6967	$\pm 0.0064 \pm 0.0109$	0.6493	$\pm 0.0061 \pm 0.0065$
0.400 - 0.440	0.5164	$\pm 0.0054 \pm 0.0092$	0.5232	$\pm 0.0055 \pm 0.0108$	0.4820	$\pm 0.0052 \pm 0.0048$
0.440 - 0.480	0.3940	$\pm 0.0047 \pm 0.0079$	0.3998	$\pm 0.0048 \pm 0.0108$	0.3493	$\pm 0.0044 \pm 0.0055$
0.480 - 0.520	0.2796	$\pm 0.0039 \pm 0.0064$	0.2788	$\pm 0.0040 \pm 0.0097$	0.2497	$\pm 0.0037 \pm 0.0065$
0.520 - 0.560	0.1835	$\pm 0.0031 \pm 0.0047$	0.1725	$\pm 0.0031 \pm 0.0076$	0.1489	$\pm 0.0028 \pm 0.0058$
0.560 - 0.600	0.1048	$\pm 0.0023 \pm 0.0030$	0.0892	$\pm 0.0022 \pm 0.0049$	0.0714	$\pm 0.0019 \pm 0.0038$
0.600 - 0.640	0.0453	$\pm 0.0015 \pm 0.0015$	0.0306	$\pm 0.0014 \pm 0.0020$	0.0203	$\pm 0.0010 \pm 0.0014$

Table 29: Major M

Minor	$\frac{1}{N} \frac{dN}{dm}$					
Interval	charged particles corrected to charged particles		charged particles corrected to charged plus neutral particles		charged plus neutral particles corrected to charged plus neutral particles	
0.000 - 0.020	0.1449	$\pm 0.0043 \pm 0.0107$	0.0689	$\pm 0.0067 \pm 0.0117$	0.0156	$\pm 0.0017 \pm 0.0036$
0.020 - 0.040	3.109	$\pm 0.019 \pm 0.031$	1.968	$\pm 0.019 \pm 0.100$	1.236	$\pm 0.013 \pm 0.066$
0.040 - 0.050	7.968	$\pm 0.044 \pm 0.080$	6.346	$\pm 0.041 \pm 0.071$	5.706	$\pm 0.037 \pm 0.073$
0.050 - 0.060	10.353	$\pm 0.050 \pm 0.104$	9.491	$\pm 0.048 \pm 0.116$	9.714	$\pm 0.048 \pm 0.125$
0.060 - 0.070	11.272	$\pm 0.052 \pm 0.113$	11.116	$\pm 0.051 \pm 0.147$	12.015	$\pm 0.054 \pm 0.155$
0.070 - 0.080	10.890	$\pm 0.051 \pm 0.109$	11.163	$\pm 0.051 \pm 0.159$	12.437	$\pm 0.055 \pm 0.161$
0.080 - 0.100	9.104	$\pm 0.033 \pm 0.091$	9.417	$\pm 0.033 \pm 0.149$	10.404	$\pm 0.036 \pm 0.136$
0.100 - 0.120	6.292	$\pm 0.027 \pm 0.063$	6.569	$\pm 0.028 \pm 0.118$	6.918	$\pm 0.029 \pm 0.092$
0.120 - 0.140	4.029	$\pm 0.022 \pm 0.048$	4.233	$\pm 0.022 \pm 0.085$	4.250	$\pm 0.023 \pm 0.058$
0.140 - 0.160	2.519	$\pm 0.017 \pm 0.037$	2.612	$\pm 0.018 \pm 0.058$	2.517	$\pm 0.017 \pm 0.035$
0.160 - 0.200	1.2832	$\pm 0.0085 \pm 0.0243$	1.3590	$\pm 0.0092 \pm 0.0349$	1.2561	$\pm 0.0086 \pm 0.0187$
0.200 - 0.240	0.5255	$\pm 0.0055 \pm 0.0130$	0.5586	$\pm 0.0059 \pm 0.0170$	0.4895	$\pm 0.0054 \pm 0.0080$
0.240 - 0.280	0.2160	$\pm 0.0035 \pm 0.0067$	0.2448	$\pm 0.0040 \pm 0.0086$	0.2112	$\pm 0.0036 \pm 0.0039$
0.280 - 0.320	0.0928	$\pm 0.0023 \pm 0.0066$	0.1020	$\pm 0.0026 \pm 0.0041$	0.0879	$\pm 0.0023 \pm 0.0018$
0.320 - 0.400	0.0248	$\pm 0.0009 \pm 0.0012$	0.0291	$\pm 0.0010 \pm 0.0014$	0.0250	$\pm 0.0009 \pm 0.0006$

Table 30: Minor m

Oblateness	$\frac{1}{N} \frac{dN}{dO}$							
Interval	charged particles corrected to charged particles			charged particles corrected to charged plus neutral particles			charged plus neutral particles corrected to charged plus neutral particles	
0.000 - 0.020	7.415	± 0.031	± 0.106	8.103	± 0.033	± 0.193	9.357	± 0.036 ± 0.178
0.020 - 0.040	10.840	± 0.037	± 0.108	10.809	± 0.036	± 0.204	11.508	± 0.038 ± 0.140
0.040 - 0.060	7.800	± 0.030	± 0.078	7.294	± 0.029	± 0.109	7.215	± 0.029 ± 0.072
0.060 - 0.080	5.273	± 0.025	± 0.053	5.030	± 0.024	± 0.060	4.736	± 0.023 ± 0.047
0.080 - 0.100	3.797	± 0.021	± 0.038	3.697	± 0.021	± 0.037	3.477	± 0.020 ± 0.035
0.100 - 0.120	2.838	± 0.018	± 0.028	2.850	± 0.018	± 0.029	2.696	± 0.018 ± 0.027
0.120 - 0.140	2.218	± 0.016	± 0.022	2.208	± 0.016	± 0.022	2.106	± 0.016 ± 0.021
0.140 - 0.160	1.760	± 0.014	± 0.018	1.793	± 0.015	± 0.018	1.690	± 0.014 ± 0.017
0.160 - 0.200	1.3199	± 0.0087	± 0.0132	1.3247	± 0.0088	± 0.0132	1.2648±	0.0085 ± 0.0126
0.200 - 0.240	0.8874	± 0.0070	± 0.0089	0.9033	± 0.0072	± 0.0091	0.8403±	0.0069 ± 0.0087
0.240 - 0.280	0.6208	± 0.0058	± 0.0078	0.6179	± 0.0060	± 0.0065	0.5674±	0.0056 ± 0.0065
0.280 - 0.320	0.4354	± 0.0049	± 0.0055	0.4268	± 0.0049	± 0.0049	0.3842±	0.0046 ± 0.0050
0.320 - 0.360	0.2959	± 0.0040	± 0.0030	0.2855	± 0.0040	± 0.0046	0.2573±	0.0037 ± 0.0043
0.360 - 0.400	0.1987	± 0.0033	± 0.0021	0.1841	± 0.0032	± 0.0045	0.1594±	0.0029 ± 0.0037
0.400 - 0.440	0.1163	± 0.0025	± 0.0032	0.0977	± 0.0022	± 0.0032	0.0836±	0.0020 ± 0.0030
0.440 - 0.520	0.0453	± 0.0011	± 0.0007	0.0293	± 0.0009	± 0.0014	0.0221±	0.0007 ± 0.0015

Table 31: Oblateness O

Heavy Jet Mass	$\frac{1}{N} \frac{dN}{dM_{high}^2/E_{vis}^2}$							
Interval	charged particles corrected to charged particles			charged particles corrected to charged plus neutral particles			charged plus neutral particles corrected to charged plus neutral particles	
0.000 - 0.010	4.450	± 0.032	± 0.260	2.035	± 0.028	± 0.180	1.994	± 0.027 ± 0.166
0.010 - 0.020	17.310	± 0.062	± 0.173	16.492	± 0.059	± 0.165	18.580	± 0.065 ± 0.709
0.020 - 0.030	17.290	± 0.066	± 0.267	20.056	± 0.074	± 0.611	20.678	± 0.076 ± 0.729
0.030 - 0.040	12.996	± 0.058	± 0.206	13.772	± 0.061	± 0.377	13.377	± 0.060 ± 0.412
0.040 - 0.050	9.496	± 0.050	± 0.154	9.456	± 0.051	± 0.233	8.965	± 0.049 ± 0.239
0.050 - 0.060	6.957	± 0.043	± 0.116	6.862	± 0.043	± 0.153	6.558	± 0.041 ± 0.151
0.060 - 0.080	4.811	± 0.025	± 0.083	4.762	± 0.025	± 0.092	4.515	± 0.024 ± 0.082
0.080 - 0.100	3.064	± 0.020	± 0.055	3.071	± 0.020	± 0.051	2.914	± 0.019 ± 0.037
0.100 - 0.120	2.109	± 0.016	± 0.039	2.126	± 0.017	± 0.032	1.991	± 0.016 ± 0.020
0.120 - 0.140	1.468	± 0.014	± 0.028	1.478	± 0.014	± 0.023	1.406	± 0.013 ± 0.014
0.140 - 0.160	1.078	± 0.012	± 0.021	1.096	± 0.012	± 0.018	1.010	± 0.011 ± 0.010
0.160 - 0.200	0.6959	± 0.0067	± 0.0142	0.6983	± 0.0068	± 0.0149	0.6319±	0.0063 ± 0.0063
0.200 - 0.250	0.3444	± 0.0042	± 0.0074	0.3461	± 0.0042	± 0.0118	0.3085±	0.0039 ± 0.0051
0.250 - 0.300	0.1497	± 0.0027	± 0.0033	0.1457	± 0.0027	± 0.0082	0.1115±	0.0022 ± 0.0039
0.300 - 0.350	0.0547	± 0.0015	± 0.0012	0.0309	± 0.0013	± 0.0027	0.0184±	0.0008 ± 0.0012
0.350 - 0.400	0.0192	± 0.0009	± 0.0004	0.0022	± 0.0006	± 0.0003	0.0008±	0.0002 ± 0.0001

Table 32: Heavy Jet Mass M_{high}^2/E_{vis}^2

Light Jet Mass			$\frac{1}{N} \frac{dN}{dM_{up}^2/E_{vis}^2}$		
Interval	charged particles corrected to charged particles		charged particles corrected to charged plus neutral particles		charged plus neutral particles corrected to charged plus neutral particles
0.000 - 0.010	39.362	± 0.093 ± 1.220	24.708	± 0.077 ± 1.403	23.414 ± 0.074 ± 1.595
0.010 - 0.020	31.177	± 0.092 ± 1.074	35.91	± 0.10 ± 2.35	39.12 ± 0.11 ± 2.65
0.020 - 0.030	13.733	± 0.065 ± 0.523	17.370	± 0.079 ± 1.196	18.080 ± 0.081 ± 1.215
0.030 - 0.040	6.486	± 0.044 ± 0.270	7.628	± 0.051 ± 0.516	7.704 ± 0.052 ± 0.514
0.040 - 0.050	3.358	± 0.032 ± 0.147	3.829	± 0.036 ± 0.251	3.922 ± 0.036 ± 0.260
0.050 - 0.060	1.879	± 0.024 ± 0.082	2.085	± 0.026 ± 0.132	2.128 ± 0.026 ± 0.140
0.060 - 0.080	0.901	± 0.012 ± 0.040	0.989	± 0.013 ± 0.063	1.013 ± 0.013 ± 0.066
0.080 - 0.100	0.3212	± 0.0072 ± 0.0175	0.3326	± 0.0073 ± 0.0234	0.3748± 0.0079 ± 0.0241
0.100 - 0.120	0.1087	± 0.0044 ± 0.0082	0.1135	± 0.0044 ± 0.0096	0.1412± 0.0050 ± 0.0089

Table 33: Light Jet Mass M_{low}^2/E_{vis}^2

Difference of the Jet Masses			$\frac{1}{N} \frac{dN}{dM_{diff}^2/E_{vis}^2}$		
Interval	charged particles corrected to charged particles		charged particles corrected to charged plus neutral particles		charged plus neutral particles corrected to charged plus neutral particles
0.000 - 0.010	29.229	$\pm 0.082 \pm 0.292$	31.955	$\pm 0.086 \pm 0.332$	35.393 $\pm 0.092 \pm 0.354$
0.010 - 0.020	19.648	$\pm 0.068 \pm 0.196$	20.327	$\pm 0.071 \pm 0.203$	20.745 $\pm 0.071 \pm 0.207$
0.020 - 0.030	12.462	$\pm 0.055 \pm 0.125$	11.909	$\pm 0.054 \pm 0.119$	11.426 $\pm 0.052 \pm 0.114$
0.030 - 0.040	8.332	$\pm 0.045 \pm 0.083$	7.654	$\pm 0.043 \pm 0.077$	7.170 $\pm 0.041 \pm 0.072$
0.040 - 0.060	5.164	$\pm 0.025 \pm 0.052$	4.706	$\pm 0.024 \pm 0.047$	4.344 $\pm 0.023 \pm 0.043$
0.060 - 0.080	2.987	$\pm 0.019 \pm 0.034$	2.816	$\pm 0.019 \pm 0.034$	2.605 $\pm 0.017 \pm 0.026$
0.080 - 0.120	1.646	$\pm 0.010 \pm 0.022$	1.5632	$\pm 0.0098 \pm 0.0267$	1.4238 $\pm 0.0092 \pm 0.0142$
0.120 - 0.160	0.8298	$\pm 0.0070 \pm 0.0136$	0.8043	$\pm 0.0070 \pm 0.0186$	0.7061 $\pm 0.0064 \pm 0.0071$
0.160 - 0.200	0.4606	$\pm 0.0053 \pm 0.0091$	0.4374	$\pm 0.0051 \pm 0.0132$	0.3831 $\pm 0.0046 \pm 0.0044$
0.200 - 0.250	0.2358	$\pm 0.0033 \pm 0.0060$	0.2223	$\pm 0.0032 \pm 0.0087$	0.1836 $\pm 0.0028 \pm 0.0032$
0.250 - 0.300	0.1043	$\pm 0.0022 \pm 0.0036$	0.0838	$\pm 0.0020 \pm 0.0042$	0.0579 $\pm 0.0015 \pm 0.0018$
0.300 - 0.350	0.0417	$\pm 0.0013 \pm 0.0020$	0.0144	$\pm 0.0010 \pm 0.0009$	0.0075 $\pm 0.0006 \pm 0.0006$
0.350 - 0.400	0.0151	$\pm 0.0008 \pm 0.0010$	0.0008	$\pm 0.0005 \pm 0.0001$	0.0003 $\pm 0.0002 \pm 0.0001$

Table 34: Difference of the Jet Masses M_{diff}^2/E_{vis}^2

Diff. Jet Rate D_2 Durham Algorithm				$\frac{1}{N} \frac{dN}{dD_2^D}$			
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles
	charged particles			charged plus neutral particles			corrected to charged plus neutral particles
0.000 - 0.010	63.52	± 0.12	± 0.64	62.64	± 0.12	± 0.63	63.81 ± 0.12 ± 0.64
0.010 - 0.020	12.230	± 0.054	± 0.122	12.326	± 0.054	± 0.126	11.589 ± 0.052 ± 0.116
0.020 - 0.030	6.128	± 0.038	± 0.061	6.324	± 0.039	± 0.080	6.040 ± 0.038 ± 0.060
0.030 - 0.040	3.828	± 0.030	± 0.038	3.993	± 0.031	± 0.058	3.823 ± 0.030 ± 0.038
0.040 - 0.050	2.756	± 0.025	± 0.030	2.888	± 0.027	± 0.046	2.697 ± 0.026 ± 0.027
0.050 - 0.060	1.995	± 0.022	± 0.028	2.095	± 0.023	± 0.036	2.041 ± 0.022 ± 0.020
0.060 - 0.080	1.470	± 0.013	± 0.025	1.484	± 0.014	± 0.027	1.410 ± 0.013 ± 0.014
0.080 - 0.100	0.932	± 0.011	± 0.019	0.963	± 0.011	± 0.020	0.952 ± 0.011 ± 0.011
0.100 - 0.120	0.6551	± 0.0091	± 0.0162	0.6537	± 0.0089	± 0.0160	0.6708 ± 0.0090 ± 0.0102
0.120 - 0.140	0.4985	± 0.0081	± 0.0144	0.4723	± 0.0076	± 0.0138	0.4831 ± 0.0077 ± 0.0098
0.140 - 0.160	0.3442	± 0.0065	± 0.0114	0.3431	± 0.0065	± 0.0120	0.3660 ± 0.0068 ± 0.0098
0.160 - 0.180	0.2619	± 0.0059	± 0.0103	0.2614	± 0.0057	± 0.0112	0.2830 ± 0.0061 ± 0.0102
0.180 - 0.200	0.2023	± 0.0055	± 0.0099	0.1841	± 0.0048	± 0.0100	0.2190 ± 0.0054 ± 0.0109
0.200 - 0.250	0.0978	± 0.0024	± 0.0060	0.1001	± 0.0023	± 0.0069	0.1357 ± 0.0028 ± 0.0092
0.250 - 0.300	0.0248	± 0.0013	± 0.0019	0.0309	± 0.0015	± 0.0027	0.0605 ± 0.0022 ± 0.0055

Table 35: Differential Jet Rate D_2 Durham Algorithm D_2^D

Diff. Jet Rate D_3 Durham Algorithm				$\frac{1}{N} \frac{dN}{dD_3^D}$			
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles
	charged particles			charged plus neutral particles			corrected to charged plus neutral particles
0.000 - 0.002	318.65	± 0.62	± 3.19	318.41	± 0.62	± 3.18	333.97 ± 0.64 ± 3.34
0.002 - 0.004	91.82	± 0.33	± 0.92	90.83	± 0.32	± 1.07	85.51 ± 0.31 ± 0.86
0.004 - 0.006	33.87	± 0.20	± 0.34	33.23	± 0.20	± 0.49	29.49 ± 0.18 ± 0.29
0.006 - 0.008	16.86	± 0.14	± 0.17	16.99	± 0.14	± 0.31	14.78 ± 0.13 ± 0.15
0.008 - 0.012	8.383	± 0.070	± 0.120	8.546	± 0.073	± 0.189	7.583 ± 0.067 ± 0.076
0.012 - 0.016	3.967	± 0.048	± 0.079	4.119	± 0.051	± 0.112	3.707 ± 0.047 ± 0.037
0.016 - 0.020	2.218	± 0.037	± 0.058	2.325	± 0.039	± 0.077	2.119 ± 0.037 ± 0.042
0.020 - 0.024	1.302	± 0.028	± 0.041	1.479	± 0.032	± 0.058	1.348 ± 0.030 ± 0.042
0.024 - 0.028	0.906	± 0.024	± 0.033	0.961	± 0.026	± 0.043	0.906 ± 0.024 ± 0.038
0.028 - 0.032	0.621	± 0.020	± 0.025	0.669	± 0.022	± 0.032	0.646 ± 0.021 ± 0.034
0.032 - 0.040	0.344	± 0.010	± 0.016	0.387	± 0.012	± 0.019	0.366 ± 0.011 ± 0.025
0.040 - 0.050	0.1892	± 0.0071	± 0.0104	0.2045	± 0.0078	± 0.0097	0.1965 ± 0.0076 ± 0.0177
0.050 - 0.060	0.0867	± 0.0050	± 0.0060	0.0869	± 0.0050	± 0.0041	0.0877 ± 0.0050 ± 0.0100
0.060 - 0.080	0.0331	± 0.0024	± 0.0030	0.0346	± 0.0023	± 0.0021	0.0367 ± 0.0024 ± 0.0054
0.080 - 0.100	0.00742	± 0.00136	± 0.00088	0.00615	± 0.00096	± 0.00066	0.0073 ± 0.0010 ± 0.0014

Table 36: Differential Jet Rate D_3 Durham Algorithm D_3^D

Diff. Jet Rate D_4 Durham Algorithm				$\frac{1}{N} \frac{dN}{dD_4^D}$					
Interval	charged particles corrected to charged particles			charged particles corrected to charged plus neutral particles			charged plus neutral particles corrected to charged plus neutral particles		
0.000 - 0.001	688.3	± 1.3	± 6.9	684.2	± 1.3	± 6.8	711.0	± 1.3	± 7.1
0.001 - 0.002	190.40	± 0.67	± 1.90	193.30	± 0.67	± 1.93	184.32	± 0.66	± 1.84
0.002 - 0.003	60.98	± 0.37	± 0.61	61.50	± 0.38	± 2.31	53.35	± 0.35	± 1.06
0.003 - 0.004	25.69	± 0.24	± 0.26	25.32	± 0.25	± 0.98	21.18	± 0.22	± 0.53
0.004 - 0.005	12.63	± 0.17	± 0.18	12.43	± 0.18	± 0.49	10.15	± 0.15	± 0.30
0.005 - 0.006	6.89	± 0.13	± 0.15	6.90	± 0.13	± 0.28	5.67	± 0.11	± 0.20
0.006 - 0.008	3.442	± 0.063	± 0.112	3.377	± 0.066	± 0.145	2.768	± 0.057	± 0.114
0.008 - 0.010	1.454	± 0.040	± 0.068	1.457	± 0.044	± 0.067	1.203	± 0.039	± 0.060
0.010 - 0.012	0.750	± 0.031	± 0.045	0.744	± 0.032	± 0.037	0.617	± 0.028	± 0.036
0.012 - 0.014	0.348	± 0.020	± 0.025	0.405	± 0.024	± 0.021	0.357	± 0.022	± 0.024
0.014 - 0.016	0.204	± 0.016	± 0.017	0.258	± 0.020	± 0.015	0.215	± 0.017	± 0.016
0.016 - 0.018	0.130	± 0.013	± 0.013	0.1551	± 0.0156	± 0.0095	0.144	± 0.015	± 0.011
0.018 - 0.020	0.0712	± 0.0092	± 0.0078	0.0901	± 0.0118	± 0.0059	0.0745	± 0.0103	± 0.0064
0.020 - 0.025	0.0342	± 0.0041	± 0.0044	0.0483	± 0.0057	± 0.0036	0.0434	± 0.0052	± 0.0042
0.025 - 0.030	0.0110	± 0.0023	± 0.0017	0.0151	± 0.0028	± 0.0014	0.0133	± 0.0028	± 0.0014

Table 37: Differential Jet Rate D_4 Durham Algorithm D_4^D

Diff. Jet Rate D_2 Jade Algorithm				$\frac{1}{N} \frac{dN}{dD_2^J}$					
Interval	charged particles corrected to charged particles			charged particles corrected to charged plus neutral particles			charged plus neutral particles corrected to charged plus neutral particles		
0.000 - 0.010	26.014	± 0.077	± 0.260	24.267	± 0.073	± 0.442	26.317	± 0.078	± 0.289
0.010 - 0.020	18.243	± 0.065	± 0.182	18.064	± 0.066	± 0.269	17.670	± 0.064	± 0.197
0.020 - 0.030	11.945	± 0.054	± 0.119	12.277	± 0.056	± 0.157	11.622	± 0.054	± 0.134
0.030 - 0.040	8.378	± 0.046	± 0.084	8.677	± 0.047	± 0.103	8.177	± 0.045	± 0.099
0.040 - 0.050	6.218	± 0.039	± 0.062	6.442	± 0.041	± 0.079	6.053	± 0.039	± 0.078
0.050 - 0.060	4.817	± 0.035	± 0.048	4.983	± 0.036	± 0.069	4.731	± 0.035	± 0.065
0.060 - 0.080	3.490	± 0.021	± 0.035	3.691	± 0.022	± 0.059	3.479	± 0.021	± 0.051
0.080 - 0.100	2.358	± 0.017	± 0.024	2.525	± 0.018	± 0.048	2.434	± 0.018	± 0.038
0.100 - 0.120	1.680	± 0.014	± 0.020	1.797	± 0.015	± 0.039	1.740	± 0.015	± 0.029
0.120 - 0.140	1.247	± 0.012	± 0.017	1.326	± 0.013	± 0.034	1.314	± 0.013	± 0.023
0.140 - 0.160	0.941	± 0.011	± 0.016	0.993	± 0.011	± 0.029	0.981	± 0.011	± 0.018
0.160 - 0.180	0.7392	± 0.0097	± 0.0154	0.7562	± 0.0098	± 0.0250	0.7495 $\pm$	± 0.0098	± 0.0150
0.180 - 0.200	0.5596	± 0.0084	± 0.0153	0.5766	± 0.0086	± 0.0207	0.5711 $\pm$	± 0.0086	± 0.0126
0.200 - 0.250	0.3330	± 0.0042	± 0.0120	0.3416	± 0.0042	± 0.0130	0.3633 $\pm$	± 0.0044	± 0.0091
0.250 - 0.300	0.1317	± 0.0028	± 0.0062	0.1296	± 0.0026	± 0.0051	0.1621 $\pm$	± 0.0030	± 0.0046

Table 38: Differential Jet Rate D_2 Jade Algorithm D_2^J

Diff. Jet Rate D_3 Jade Algorithm				$\frac{1}{N} \frac{dN}{dD_3^J}$					
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles corrected to		
	charged particles			charged plus neutral particles			charged plus neutral particles		
0.000 - 0.002	107.87	± 0.36	± 1.08	103.55	± 0.35	± 1.04	113.89	± 0.37	± 1.14
0.002 - 0.004	121.68	± 0.38	± 1.22	122.74	± 0.37	± 1.41	130.22	± 0.39	± 1.30
0.004 - 0.006	74.79	± 0.29	± 0.75	73.19	± 0.29	± 0.95	71.16	± 0.28	± 0.80
0.006 - 0.008	46.84	± 0.23	± 0.47	47.04	± 0.24	± 0.69	43.53	± 0.22	± 0.51
0.008 - 0.012	27.71	± 0.13	± 0.28	28.13	± 0.13	± 0.48	25.56	± 0.12	± 0.32
0.012 - 0.016	15.164	± 0.097	± 0.152	15.73	± 0.10	± 0.32	14.035	± 0.095	± 0.191
0.016 - 0.020	9.124	± 0.075	± 0.091	9.798	± 0.082	± 0.228	8.816	± 0.075	± 0.133
0.020 - 0.024	5.939	± 0.060	± 0.070	6.363	± 0.066	± 0.168	5.733	± 0.061	± 0.095
0.024 - 0.028	4.101	± 0.051	± 0.064	4.417	± 0.056	± 0.130	4.009	± 0.051	± 0.074
0.028 - 0.032	2.920	± 0.043	± 0.057	3.163	± 0.047	± 0.103	2.897	± 0.044	± 0.059
0.032 - 0.040	1.857	± 0.024	± 0.047	2.026	± 0.027	± 0.076	1.849	± 0.025	± 0.044
0.040 - 0.050	0.970	± 0.016	± 0.033	1.058	± 0.018	± 0.047	0.988	± 0.017	± 0.029
0.050 - 0.060	0.509	± 0.012	± 0.022	0.542	± 0.013	± 0.028	0.513	± 0.012	± 0.019
0.060 - 0.080	0.1952	± 0.0051	± 0.0113	0.2241	± 0.0057	± 0.0144	0.2203	± 0.0057	± 0.0114
0.080 - 0.100	0.0554	± 0.0029	± 0.0043	0.0594	± 0.0031	± 0.0047	0.0654	± 0.0033	± 0.0049

Table 39: Differential Jet Rate D_3 Jade Algorithm D_3^J

Diff. Jet Rate D_4 Jade Algorithm				$\frac{1}{N} \frac{dN}{dD_4^J}$					
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles corrected to		
	charged particles			charged plus neutral particles			charged plus neutral particles		
0.000 - 0.001	248.15	± 0.77	± 2.48	227.18	± 0.75	± 3.97	240.63	± 0.78	± 4.58
0.001 - 0.002	286.76	± 0.82	± 2.87	289.93	± 0.82	± 5.54	309.26	± 0.86	± 4.96
0.002 - 0.003	167.03	± 0.63	± 1.67	173.10	± 0.64	± 3.59	171.69	± 0.64	± 2.39
0.003 - 0.004	97.00	± 0.48	± 1.04	99.50	± 0.49	± 2.22	92.65	± 0.47	± 1.18
0.004 - 0.005	59.58	± 0.38	± 0.71	60.08	± 0.39	± 1.44	52.87	± 0.35	± 0.65
0.005 - 0.006	37.48	± 0.30	± 0.49	38.54	± 0.31	± 0.98	33.41	± 0.28	± 0.43
0.006 - 0.008	21.61	± 0.16	± 0.32	22.33	± 0.17	± 0.62	18.83	± 0.15	± 0.27
0.008 - 0.010	11.10	± 0.12	± 0.19	11.66	± 0.13	± 0.36	9.45	± 0.11	± 0.17
0.010 - 0.012	6.235	± 0.087	± 0.124	6.717	± 0.098	± 0.231	5.567	± 0.086	± 0.122
0.012 - 0.014	3.602	± 0.065	± 0.080	4.068	± 0.077	± 0.153	3.262	± 0.066	± 0.089
0.014 - 0.016	2.205	± 0.052	± 0.055	2.454	± 0.060	± 0.100	2.047	± 0.052	± 0.070
0.016 - 0.018	1.373	± 0.041	± 0.037	1.556	± 0.048	± 0.069	1.330	± 0.042	± 0.055
0.018 - 0.020	0.874	± 0.032	± 0.026	1.035	± 0.040	± 0.049	0.912	± 0.036	± 0.044
0.020 - 0.025	0.460	± 0.015	± 0.016	0.538	± 0.018	± 0.028	0.456	± 0.016	± 0.025
0.025 - 0.030	0.1735	± 0.0096	± 0.0070	0.202	± 0.011	± 0.012	0.195	± 0.011	± 0.012
0.030 - 0.050	0.0303	± 0.0021	± 0.0017	0.037	± 0.002	± 0.003	0.034	± 0.003	± 0.002

Table 40: Differential Jet Rate D_4 Jade Algorithm D_4^J

Wide Jet Broadening				$\frac{1}{N} \frac{dN}{dB_W}$			
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles
	charged particles			charged plus neutral particles			corrected to charged plus neutral particles
0.010 - 0.020	2.027	± 0.023	± 0.020	1.396	± 0.023	± 0.014	0.6707 $\pm 0.0096 \pm 0.1077$
0.020 - 0.030	9.923	± 0.051	± 0.232	9.815	± 0.051	± 0.747	7.538 $\pm 0.038 \pm 0.809$
0.030 - 0.040	14.758	± 0.061	± 0.245	15.520	± 0.062	± 0.819	14.690 $\pm 0.058 \pm 0.745$
0.040 - 0.050	13.374	± 0.056	± 0.140	12.966	± 0.054	± 0.413	13.942 $\pm 0.057 \pm 0.592$
0.050 - 0.060	10.620	± 0.050	± 0.106	10.186	± 0.048	± 0.136	11.298 $\pm 0.053 \pm 0.379$
0.060 - 0.070	8.369	± 0.044	± 0.084	8.211	± 0.044	± 0.082	9.065 $\pm 0.048 \pm 0.266$
0.070 - 0.080	6.811	± 0.040	± 0.068	6.790	± 0.040	± 0.068	7.387 $\pm 0.043 \pm 0.222$
0.080 - 0.100	5.146	± 0.025	± 0.051	5.166	± 0.025	± 0.052	5.445 $\pm 0.026 \pm 0.176$
0.100 - 0.120	3.584	± 0.020	± 0.036	3.685	± 0.021	± 0.037	3.796 $\pm 0.022 \pm 0.127$
0.120 - 0.140	2.577	± 0.017	± 0.026	2.651	± 0.018	± 0.027	2.670 $\pm 0.018 \pm 0.087$
0.140 - 0.170	1.733	± 0.012	± 0.017	1.780	± 0.012	± 0.018	1.756 $\pm 0.012 \pm 0.051$
0.170 - 0.200	1.0540	± 0.0090	± 0.0105	1.0990	± 0.0094	± 0.0110	1.0580 $\pm 0.0092 \pm 0.0218$
0.200 - 0.240	0.5388	± 0.0056	± 0.0135	0.5548	± 0.0058	± 0.0321	0.5288 $\pm 0.0056 \pm 0.0053$
0.240 - 0.280	0.1710	± 0.0031	± 0.0104	0.1687	± 0.0031	± 0.0284	0.1460 $\pm 0.0028 \pm 0.0071$
0.280 - 0.320	0.0262	± 0.0011	± 0.0028	0.0057	± 0.0008	± 0.0018	0.0029 $\pm 0.0004 \pm 0.0003$

Table 41: Wide Jet Broadening B_W

Small Jet Broadening				$\frac{1}{N} \frac{dN}{dB_{min}}$			
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles
	charged particles			charged plus neutral particles			corrected to charged plus neutral particles
0.000 - 0.010	2.502	± 0.025	± 0.025	1.552	± 0.026	± 0.115	0.645 $\pm 0.010 \pm 0.096$
0.010 - 0.020	18.823	± 0.068	± 0.188	15.484	± 0.064	± 0.243	11.169 $\pm 0.045 \pm 1.006$
0.020 - 0.030	28.683	± 0.084	± 0.287	29.092	± 0.084	± 0.291	28.908 $\pm 0.082 \pm 1.823$
0.030 - 0.040	20.703	± 0.071	± 0.207	22.082	± 0.073	± 0.275	25.972 $\pm 0.083 \pm 1.478$
0.040 - 0.050	11.654	± 0.052	± 0.117	11.600	± 0.052	± 0.139	14.119 $\pm 0.061 \pm 0.860$
0.050 - 0.060	6.558	± 0.039	± 0.066	6.423	± 0.038	± 0.089	7.500 $\pm 0.044 \pm 0.494$
0.060 - 0.080	3.191	± 0.019	± 0.032	3.205	± 0.019	± 0.052	3.405 $\pm 0.021 \pm 0.233$
0.080 - 0.100	1.295	± 0.012	± 0.026	1.376	± 0.013	± 0.034	1.320 $\pm 0.013 \pm 0.089$
0.100 - 0.120	0.5561	± 0.0078	± 0.0185	0.5970	± 0.0086	± 0.0222	0.5448 $\pm 0.0082 \pm 0.0328$
0.120 - 0.150	0.1921	± 0.0037	± 0.0054	0.2114	± 0.0042	± 0.0070	0.1916 $\pm 0.0040 \pm 0.0104$
0.150 - 0.180	0.0319	± 0.0015	± 0.0012	0.0356	± 0.0017	± 0.0016	0.0366 $\pm 0.0017 \pm 0.0034$

Table 42: Small Jet Broadening B_{min}

Total Jet Broadening				$\frac{1}{N} \frac{dN}{dB_{sum}}$			
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles
	charged particles			charged plus neutral particles			corrected to charged plus neutral particles
0.020 - 0.030	0.956	± 0.016	± 0.010	0.550	± 0.017	± 0.030	0.2030 $\pm 0.0055 \pm 0.0383$
0.030 - 0.040	3.765	± 0.031	± 0.038	2.804	± 0.029	± 0.033	1.628 $\pm 0.015 \pm 0.183$
0.040 - 0.050	7.117	± 0.043	± 0.071	6.646	± 0.042	± 0.066	4.999 $\pm 0.031 \pm 0.463$
0.050 - 0.060	9.301	± 0.048	± 0.093	9.342	± 0.048	± 0.093	8.190 $\pm 0.041 \pm 0.644$
0.060 - 0.070	10.059	± 0.050	± 0.101	10.265	± 0.050	± 0.103	9.887 $\pm 0.047 \pm 0.661$
0.070 - 0.080	9.212	± 0.047	± 0.092	9.272	± 0.047	± 0.093	9.883 $\pm 0.049 \pm 0.564$
0.080 - 0.090	7.958	± 0.043	± 0.080	7.915	± 0.043	± 0.079	9.007 $\pm 0.047 \pm 0.443$
0.090 - 0.100	6.815	± 0.040	± 0.068	6.788	± 0.040	± 0.068	7.746 $\pm 0.044 \pm 0.332$
0.100 - 0.110	5.898	± 0.037	± 0.059	5.899	± 0.037	± 0.059	6.714 $\pm 0.041 \pm 0.255$
0.110 - 0.130	4.785	± 0.024	± 0.048	4.812	± 0.024	± 0.048	5.393 $\pm 0.026 \pm 0.180$
0.130 - 0.150	3.619	± 0.021	± 0.036	3.667	± 0.021	± 0.038	3.998 $\pm 0.023 \pm 0.125$
0.150 - 0.170	2.737	± 0.018	± 0.027	2.799	± 0.018	± 0.033	2.980 $\pm 0.019 \pm 0.098$
0.170 - 0.190	2.143	± 0.016	± 0.021	2.203	± 0.016	± 0.031	2.294 $\pm 0.017 \pm 0.085$
0.190 - 0.210	1.670	± 0.014	± 0.017	1.728	± 0.014	± 0.029	1.747 $\pm 0.015 \pm 0.075$
0.210 - 0.240	1.2151	± 0.0096	± 0.0122	1.255	± 0.010	± 0.024	1.242 $\pm 0.010 \pm 0.063$
0.240 - 0.270	0.8189	± 0.0078	± 0.0082	0.8504	± 0.0082	± 0.0194	0.8125 $\pm 0.0080 \pm 0.0469$
0.270 - 0.300	0.5167	± 0.0061	± 0.0065	0.5243	± 0.0063	± 0.0141	0.4974 $\pm 0.0062 \pm 0.0296$
0.300 - 0.330	0.2701	± 0.0044	± 0.0054	0.2599	± 0.0045	± 0.0087	0.2285 $\pm 0.0041 \pm 0.0119$
0.330 - 0.360	0.0856	± 0.0024	± 0.0028	0.0831	± 0.0027	± 0.0042	0.0732 $\pm 0.0024 \pm 0.0007$

Table 43: Total Jet Broadening B_{sum}

Difference of the Jet Broadenings				$\frac{1}{N} \frac{dN}{dB_{diff}}$			
Interval	charged particles corrected to			charged particles corrected to			charged plus neutral particles
	charged particles			charged plus neutral particles			corrected to charged plus neutral particles
0.000 - 0.010	24.778	± 0.078	± 0.248	26.513	± 0.082	± 0.265	26.630 $\pm 0.081 \pm 0.459$
0.010 - 0.020	18.957	± 0.068	± 0.190	18.385	± 0.066	± 0.184	18.684 $\pm 0.066 \pm 0.292$
0.020 - 0.030	13.050	± 0.056	± 0.131	12.151	± 0.053	± 0.122	12.343 $\pm 0.054 \pm 0.186$
0.030 - 0.040	9.129	± 0.046	± 0.091	8.722	± 0.045	± 0.087	8.819 $\pm 0.046 \pm 0.134$
0.040 - 0.050	6.689	± 0.040	± 0.067	6.562	± 0.039	± 0.066	6.688 $\pm 0.040 \pm 0.106$
0.050 - 0.060	5.220	± 0.035	± 0.052	5.134	± 0.035	± 0.051	5.111 $\pm 0.035 \pm 0.084$
0.060 - 0.070	4.087	± 0.031	± 0.041	4.083	± 0.031	± 0.043	4.071 $\pm 0.031 \pm 0.068$
0.070 - 0.080	3.314	± 0.028	± 0.033	3.293	± 0.028	± 0.043	3.271 $\pm 0.028 \pm 0.054$
0.080 - 0.090	2.677	± 0.025	± 0.027	2.674	± 0.025	± 0.042	2.681 $\pm 0.025 \pm 0.043$
0.090 - 0.100	2.218	± 0.023	± 0.022	2.240	± 0.023	± 0.040	2.233 $\pm 0.023 \pm 0.035$
0.100 - 0.120	1.662	± 0.014	± 0.017	1.688	± 0.014	± 0.033	1.647 $\pm 0.014 \pm 0.026$
0.120 - 0.140	1.145	± 0.012	± 0.011	1.162	± 0.012	± 0.025	1.111 $\pm 0.011 \pm 0.019$
0.140 - 0.160	0.8022	± 0.0098	± 0.0108	0.8134	± 0.0099	± 0.0218	0.7618 $\pm 0.0095 \pm 0.0144$
0.160 - 0.180	0.5135	± 0.0078	± 0.0109	0.5384	± 0.0081	± 0.0181	0.5138 $\pm 0.0078 \pm 0.0119$
0.180 - 0.200	0.3325	± 0.0063	± 0.0091	0.3481	± 0.0067	± 0.0143	0.3167 $\pm 0.0062 \pm 0.0098$
0.200 - 0.240	0.1540	± 0.0029	± 0.0053	0.1605	± 0.0031	± 0.0097	0.1265 $\pm 0.0026 \pm 0.0056$
0.240 - 0.280	0.0389	± 0.0014	± 0.0023	0.0206	± 0.0013	± 0.0025	0.0117 $\pm 0.0008 \pm 0.0008$

Table 44: Difference of the Jet Broadenings B_{diff} .

C-Parameter	$\frac{1}{N} \frac{dN}{dC}$		
Interval	charged particles corrected to charged particles	charged particles corrected to charged plus neutral particles	charged plus neutral particles corrected to charged plus neutral particles
0.000 - 0.040	0.3806 ± 0.0047 ± 0.0140	0.1375 ± 0.0047 ± 0.0138	0.0881± 0.0030 ± 0.0067
0.040 - 0.080	2.388 ± 0.012 ± 0.050	1.5046 ± 0.0098 ± 0.0858	1.5383± 0.0100 ± 0.0831
0.080 - 0.120	3.653 ± 0.015 ± 0.042	3.514 ± 0.014 ± 0.108	3.909 ± 0.016 ± 0.142
0.120 - 0.160	3.323 ± 0.014 ± 0.033	3.570 ± 0.015 ± 0.058	3.833 ± 0.016 ± 0.088
0.160 - 0.200	2.637 ± 0.013 ± 0.026	2.748 ± 0.013 ± 0.027	2.835 ± 0.013 ± 0.040
0.200 - 0.240	2.091 ± 0.011 ± 0.021	2.149 ± 0.011 ± 0.021	2.164 ± 0.012 ± 0.022
0.240 - 0.280	1.680 ± 0.010 ± 0.017	1.732 ± 0.010 ± 0.017	1.716 ± 0.010 ± 0.017
0.280 - 0.320	1.3757 ± 0.0092 ± 0.0138	1.4127 ± 0.0093 ± 0.0141	1.3860± 0.0092 ± 0.0139
0.320 - 0.360	1.1545 ± 0.0083 ± 0.0129	1.1920 ± 0.0086 ± 0.0122	1.1623± 0.0084 ± 0.0116
0.360 - 0.400	0.9709 ± 0.0076 ± 0.0127	1.0093 ± 0.0079 ± 0.0115	0.9720± 0.0077 ± 0.0097
0.400 - 0.440	0.8100 ± 0.0068 ± 0.0121	0.8629 ± 0.0074 ± 0.0108	0.8349± 0.0072 ± 0.0083
0.440 - 0.480	0.7162 ± 0.0065 ± 0.0120	0.7545 ± 0.0068 ± 0.0105	0.7161± 0.0066 ± 0.0072
0.480 - 0.520	0.6247 ± 0.0060 ± 0.0116	0.6544 ± 0.0064 ± 0.0101	0.6205± 0.0061 ± 0.0062
0.520 - 0.560	0.5573 ± 0.0057 ± 0.0113	0.5914 ± 0.0061 ± 0.0103	0.5441± 0.0057 ± 0.0054
0.560 - 0.600	0.4939 ± 0.0053 ± 0.0108	0.5187 ± 0.0057 ± 0.0101	0.4844± 0.0054 ± 0.0050
0.600 - 0.640	0.4245 ± 0.0049 ± 0.0100	0.4558 ± 0.0053 ± 0.0101	0.4209± 0.0050 ± 0.0063
0.640 - 0.680	0.3860 ± 0.0048 ± 0.0097	0.3968 ± 0.0049 ± 0.0101	0.3699± 0.0046 ± 0.0079
0.680 - 0.720	0.3589 ± 0.0046 ± 0.0095	0.3697 ± 0.0047 ± 0.0106	0.3286± 0.0044 ± 0.0099
0.720 - 0.760	0.3063 ± 0.0041 ± 0.0084	0.3250 ± 0.0044 ± 0.0104	0.2813± 0.0040 ± 0.0129
0.760 - 0.800	0.2672 ± 0.0039 ± 0.0073	0.2807 ± 0.0042 ± 0.0096	0.2178± 0.0033 ± 0.0151
0.800 - 0.840	0.1485 ± 0.0028 ± 0.0037	0.1803 ± 0.0035 ± 0.0062	0.1287± 0.0026 ± 0.0130
0.840 - 0.880	0.0716 ± 0.0020 ± 0.0015	0.0831 ± 0.0023 ± 0.0026	0.0542± 0.0016 ± 0.0076
0.880 - 0.920	0.0307 ± 0.0013 ± 0.0004	0.0361 ± 0.0016 ± 0.0008	0.0212± 0.0009 ± 0.0040

Table 45: C-Parameter C

D-Parameter	$\frac{1}{N} \frac{dN}{dD}$		
Interval	charged particles corrected to charged particles	charged particles corrected to charged plus neutral particles	charged plus neutral particles corrected to charged plus neutral particles
0.000 - 0.008	29.731 ± 0.093 ± 0.308	22.567 ± 0.083 ± 0.645	22.228 ± 0.082 ± 0.868
0.008 - 0.016	19.922 ± 0.077 ± 0.199	20.963 ± 0.080 ± 0.401	22.766 ± 0.085 ± 0.440
0.016 - 0.030	11.026 ± 0.044 ± 0.110	11.522 ± 0.045 ± 0.169	12.107 ± 0.047 ± 0.150
0.030 - 0.044	6.513 ± 0.034 ± 0.065	6.763 ± 0.035 ± 0.093	6.879 ± 0.035 ± 0.079
0.044 - 0.066	4.066 ± 0.021 ± 0.045	4.262 ± 0.022 ± 0.064	4.284 ± 0.022 ± 0.053
0.066 - 0.088	2.640 ± 0.017 ± 0.033	2.790 ± 0.018 ± 0.048	2.727 ± 0.018 ± 0.036
0.088 - 0.112	1.864 ± 0.014 ± 0.025	1.964 ± 0.014 ± 0.039	1.909 ± 0.014 ± 0.028
0.112 - 0.136	1.352 ± 0.012 ± 0.019	1.469 ± 0.013 ± 0.033	1.415 ± 0.012 ± 0.022
0.136 - 0.162	1.0013 ± 0.0095 ± 0.0149	1.113 ± 0.011 ± 0.028	1.051 ± 0.010 ± 0.018
0.162 - 0.188	0.7829 ± 0.0085 ± 0.0123	0.8654 ± 0.0094 ± 0.0239	0.7977± 0.0089 ± 0.0145
0.188 - 0.218	0.6187 ± 0.0071 ± 0.0103	0.6835 ± 0.0078 ± 0.0204	0.6155± 0.0073 ± 0.0117
0.218 - 0.248	0.4490 ± 0.0059 ± 0.0081	0.5253 ± 0.0069 ± 0.0168	0.4566± 0.0063 ± 0.0089
0.248 - 0.284	0.3425 ± 0.0048 ± 0.0068	0.3915 ± 0.0055 ± 0.0134	0.3341± 0.0049 ± 0.0065
0.284 - 0.320	0.2575 ± 0.0042 ± 0.0057	0.2834 ± 0.0046 ± 0.0103	0.2452± 0.0042 ± 0.0049
0.320 - 0.360	0.1884 ± 0.0034 ± 0.0047	0.2142 ± 0.0039 ± 0.0083	0.1774± 0.0033 ± 0.0037
0.360 - 0.400	0.1363 ± 0.0028 ± 0.0038	0.1629 ± 0.0034 ± 0.0067	0.1234± 0.0028 ± 0.0028
0.400 - 0.450	0.0962 ± 0.0021 ± 0.0029	0.1095 ± 0.0025 ± 0.0049	0.0902± 0.0021 ± 0.0023
0.450 - 0.500	0.0644 ± 0.0017 ± 0.0021	0.0758 ± 0.0020 ± 0.0036	0.0603± 0.0017 ± 0.0018
0.500 - 0.560	0.0394 ± 0.0012 ± 0.0014	0.0466 ± 0.0015 ± 0.0023	0.0368± 0.0012 ± 0.0013
0.560 - 0.620	0.0277 ± 0.0010 ± 0.0010	0.0314 ± 0.0012 ± 0.0016	0.0222± 0.0009 ± 0.0009
0.620 - 0.710	0.0151 ± 0.0006 ± 0.0006	0.0182 ± 0.0007 ± 0.0009	0.0128± 0.0006 ± 0.0006
0.710 - 0.800	0.0071 ± 0.0004 ± 0.0003	0.0084 ± 0.0005 ± 0.0040	0.0052± 0.0004 ± 0.0003

Table 46: D-Parameter D

E Tables of Results of the Fits

Parameter	Default	Range gen.	Fit Result		
			Val.	stat.	sys.
QCDLAM	0.18	0.155 - 0.205	0.163±0.001	$^{+0.004}_{-0.005}$	
RMASS(13)	0.75	0.64 - 0.96	0.65 ±0.01	$^{+0.02}_{-0.01}$	
CLMAX	3.35	2.35 - 4.35	3.48 ±0.04	$^{+0.20}_{-0.13}$	
CLPOW	2.0	1.0 - 2.0	1.49 ±0.04	$^{+0.16}_{-0.08}$	
CLSMR	0.	0.0 - 1.0	0.36 ±0.04	$^{+0.06}_{-0.01}$	
DECWT	1.0	0.0 - 1.0	0.77 ±0.08	$^{+0.28}_{-0.24}$	
PWT(3)	1.0	0.0 - 1.0	0.83 ±0.02	$^{+0.03}_{-0.09}$	
PWT(7)	1.0	0.0 - 1.0	0.74 ±0.09	$^{+0.37}_{-0.25}$	

Table 47: Parameter setting and fit results for HERWIG 5.8 C

Parameter	Name	Default	Range gen.	Fit Result		
				Val.	stat.	sys.
λ_{QCD}	PARJ(81)	0.4	0.25 - 0.35	0.297	± 0.005	$^{+0.012}_{-0.007}$
Q_0	PARJ(82)	1.0	1.0 - 2.0	1.34	± 0.08	$^{+0.15}_{-0.09}$
a	PARJ(41)	0.5	0.1 - 0.5	0.409	± 0.020	$^{+0.009}_{-0.013}$
b	PARJ(42)	0.9	0.844	optimized		
σ_q	PARJ(21)	0.35	0.36 - 0.44	0.401	± 0.006	$^{+0.003}_{-0.013}$
$P(^1S_0)_{ud}$	-	0.5	0.3 - 0.5	0.272	± 0.030	$^{+0.108}_{-0.008}$
$P(^3S_1)_{ud}$	-	0.5	0.2 - 0.4	0.306	± 0.049	$^{+0.006}_{-0.031}$
$P(^1P_1)_{ud}$	-	0.	see text	0.094		
$P(oth.P - states)_{ud}$	-	0.	see text	0.305		
γ_s	PARJ(2)	0.30	0.27 - 0.31	0.307	± 0.007	$^{+0.002}_{-0.017}$
$P(^1S_0)_s$	-	0.4	0.3 - 0.5	0.416	± 0.029	$^{+0.015}_{-0.003}$
$P(^3S_1)_s$	-	0.6	0.2 - 0.4	0.304	± 0.023	$^{+0.017}_{-0.002}$
$P(P - states)_s$	-	0.	see text	0.280		
ϵ_c	PARJ(54)	-	variable	-0.0372	± 0.0008	$^{+0.0020}_{-0.0012}$
$P(^1S_0)_c$	-	0.25	0.26	adj. to data		
$P(^3S_1)_c$	-	0.75	0.44			
$P(P - states)_c$	-	0.	0.3			
ϵ_b	PARJ(55)	-	variable	-0.00284	± 0.00006	$^{+0.00013}_{-0.00009}$
$P(^1S_0)_b$	-	0.25	0.175	adj. to data		
$P(^3S_1)_b$	-	0.75	0.525			
$P(P - states)_b$	-	0.	0.3			
$P(qq)/P(q)$	PARJ(1)	0.1	0.08 - 0.11	0.095	± 0.001	$^{+0.004}_{-0.002}$
$P(us)/P(ud)/\gamma_s$	PARJ(3)	0.4	0.65	adj. to data		
$P(ud1)/P(ud0)$	PARJ(4)	0.05	0.07	adj. to data		
extra baryon supp.	PARJ(19)	0.	0.5	adj. to data only uds		
extra η supp.	PARJ(25)	1.0	0.65	0.65 ± 0.06		
extra η' supp.	PARJ(26)	1.0	0.23	0.23 ± 0.05		

Table 48: Parameter setting and fit results for JETSET 7.3 PS with DELPHI decays

Parameter	Name	Default	Range gen.	Fit Result		
				Val.	stat.	sys.
λ_{QCD}	PARJ(81)	0.4	0.25 - 0.35	0.297	± 0.004	$^{+0.007}_{-0.008}$
Q_0	PARJ(82)	1.0	1.0 - 2.0	1.56	± 0.11	$^{+0.21}_{-0.15}$
a	PARJ(41)	0.5	0.1 - 0.5	0.417	± 0.022	$^{+0.011}_{-0.015}$
b	PARJ(42)	0.9	0.850	optimized		
σ_q	PARJ(21)	0.35	0.36 - 0.44	0.408	± 0.005	$^{+0.004}_{-0.004}$
$P(^1S_0)_{ud}$	-	0.5	0.3 - 0.5	0.297	± 0.021	$^{+0.102}_{-0.011}$
$P(^3S_1)_{ud}$	-	0.5	0.2 - 0.4	0.289	± 0.038	$^{+0.004}_{-0.026}$
$P(^1P_1)_{ud}$	-	0.	see text	0.096		
$P(oth.P - states)_{ud}$	-	0.	see text	0.318		
γ_s	PARJ(2)	0.30	0.27 - 0.31	0.308	± 0.007	$^{+0.004}_{-0.036}$
$P(^1S_0)_s$	-	0.4	0.3 - 0.5	0.410	± 0.038	$^{+0.026}_{-0.013}$
$P(^3S_1)_s$	-	0.6	0.2 - 0.4	0.297	± 0.021	$^{+0.020}_{-0.004}$
$P(P - states)_s$	-	0.	see text	0.293		
ϵ_c	PARJ(54)	-	variable	-0.0372	± 0.0007	$^{+0.0011}_{-0.0012}$
$P(^1S_0)_c$	-	0.25	0.26	adj. to data		
$P(^3S_1)_c$	-	0.75	0.44			
$P(P - states)_c$	-	0.	0.3			
ϵ_b	PARJ(55)	-	variable	-0.00284	± 0.00005	$^{+0.00012}_{-0.00010}$
$P(^1S_0)_b$	-	0.25	0.175	adj. to data		
$P(^3S_1)_b$	-	0.75	0.525			
$P(P - states)_b$	-	0.	0.3			
$P(qq)/P(q)$	PARJ(1)	0.1	0.08 - 0.11	0.099	± 0.001	$^{+0.005}_{-0.002}$
$P(us)/P(ud)/\gamma_s$	PARJ(3)	0.4	0.65	adj. to data		
$P(ud1)/P(ud0)$	PARJ(4)	0.05	0.07	adj. to data		
extra baryon supp.	PARJ(19)	0.	0.5	adj. to data only uds		
extra η supp.	PARJ(25)	1.0	0.65	0.65 ± 0.06		
extra η' supp.	PARJ(26)	1.0	0.23	0.23 ± 0.05		

Table 49: Parameter setting and fit results for JETSET 7.4 PS with default decays

Parameter	Name	Default	Range gen.	Fit Result		
				Val.	stat.	sys.
λ_{QCD}	PARA(1)	0.22	0.20 - 0.30	0.237	± 0.003	$^{+0.006}_{-0.002}$
p_t^{QCD}	PARA(3)	0.6	0.4 - 1.2	0.64	± 0.08	$^{+0.20}_{-0.08}$
a	PARJ(41)	0.5	0.1 - 0.5	0.391	± 0.047	$^{+0.039}_{-0.046}$
b	PARJ(42)	0.9	0.850	optimized		
σ_q	PARJ(21)	0.35	0.36 - 0.44	0.390	± 0.005	$^{+0.003}_{-0.007}$
$P(^1S_0)_{ud}$	-	0.5	0.3 - 0.5	0.276	± 0.017	$^{+0.103}_{-0.012}$
$P(^3S_1)_{ud}$	-	0.5	0.2 - 0.4	0.292	± 0.056	$^{+0.003}_{-0.020}$
$P(^1P_1)_{ud}$	-	0.	see text	0.096		
$P(oth.P - states)_{ud}$	-	0.	see text	0.336		
γ_s	PARJ(2)	0.30	0.27 - 0.31	0.302	± 0.004	$^{+0.004}_{-0.008}$
$P(^1S_0)_s$	-	0.4	0.3 - 0.5	0.389	± 0.026	$^{+0.012}_{-0.010}$
$P(^3S_1)_s$	-	0.6	0.2 - 0.4	0.301	± 0.019	$^{+0.004}_{-0.002}$
$P(P - states)_s$	-	0.	see text	0.310		
ϵ_c	PARJ(54)	-	variable	-0.0378	± 0.0005	$^{+0.0009}_{-0.0004}$
$P(^1S_0)_c$	-	0.25	0.26	adj. to data		
$P(^3S_1)_c$	-	0.75	0.44			
$P(P - states)_c$	-	0.	0.3			
ϵ_b	PARJ(55)	-	variable	-0.00255	± 0.00004	$^{+0.00006}_{-0.00002}$
$P(^1S_0)_b$	-	0.25	0.175	adj. to data		
$P(^3S_1)_b$	-	0.75	0.525			
$P(P - states)_b$	-	0.	0.3			
$P(qq)/P(q)$	PARJ(1)	0.1	0.08 - 0.11	0.096	± 0.001	$^{+0.008}_{-0.001}$
$P(us)/P(ud)/\gamma_s$	PARJ(3)	0.4	0.65	adj. to data		
$P(ud1)/P(ud0)$	PARJ(4)	0.05	0.07	adj. to data		
extra baryon supp.	PARJ(19)	0.	0.5	adj. to data only uds		
extra η supp.	PARJ(25)	1.0	0.65	0.65 ± 0.06		
extra η' supp.	PARJ(26)	1.0	0.23	0.23 ± 0.05		

Table 50: Parameter setting and fit results for ARIADNE 4.06 with DELPHI decays

Parameter	name	Default	gen. range	Results		
				Value	stat.	sys.
λ_{QCD}	PARA(1)	0.22	0.20 - 0.30	0.237	± 0.004	$^{+0.005}_{-0.004}$
p_t^{QCD}	PARA(3)	0.6	0.4 - 1.2	0.68	± 0.06	$^{+0.11}_{-0.11}$
a	PARJ(41)	0.5	0.1 - 0.5	0.388	± 0.030	$^{+0.036}_{-0.112}$
b	PARJ(42)	0.9	0.850	optimized		
σ_q	PARJ(21)	0.35	0.36 - 0.44	0.387	± 0.008	$^{+0.012}_{-0.016}$
$P(^1S_0)_{ud}$	-	0.5	0.3 - 0.5	0.326	± 0.017	$^{+0.058}_{-0.023}$
$P(^3S_1)_{ud}$	-	0.5	0.2 - 0.4	0.263	± 0.039	$^{+0.015}_{-0.010}$
$P(^1P_1)_{ud}$	-	0.	see text	0.096		
$P(oth.P - states)_{ud}$	-	0.	see tex	0.411		
γ_s	PARJ(2)	0.30	0.27 - 0.31	0.282	± 0.005	$^{+0.015}_{-0.004}$
$P(^1S_0)_s$	-	0.4	0.3 - 0.5	0.395	± 0.023	$^{+0.008}_{-0.040}$
$P(^3S_1)_s$	-	0.6	0.2 - 0.4	0.325	± 0.015	$^{+0.005}_{-0.033}$
$P(P - states)_s$	-	0.	siehe TEXT	0.280		
ϵ_c	PARJ(54)	-	variable	-0.0378	± 0.0005	$^{+0.0009}_{-0.0004}$
$P(^1S_0)_c$	-	0.25	0.26	adjusted		
$P(^3S_1)_c$	-	0.75	0.44			
$P(P - states)_c$	-	0.	0.3			
ϵ_b	PARJ(55)	-	variable	-0.00255	± 0.00004	$^{+0.00006}_{-0.00002}$
$P(^1S_0)_b$	-	0.25	0.175	adjusted		
$P(^3S_1)_b$	-	0.75	0.525			
$P(P - states)_b$	-	0.	0.3			
$P(qq)/P(q)$	PARJ(1)	0.1	0.08 - 0.11	0.101	± 0.001	$^{+0.006}_{-0.001}$
$P(us)/P(ud)/\gamma_s$	PARJ(3)	0.4	0.65	adjusted		
$P(ud1)/P(ud0)$	PARJ(4)	0.05	0.07	adjusted		
extra baryon supp.	PARJ(19)	0.	0.5	adjusted only uds		
extra η supp.	PARJ(25)	1.0	0.65	0.65 ± 0.06		
extra η' supp.	PARJ(26)	1.0	0.23	0.23 ± 0.05		

Table 51: Parameter setting and fit results for ARIADNE 4.06 with default decays

Parameter	Name	Default	Range gen.		Fit Result		
					Val.	stat.	sys.
λ_{QCD}	PARJ(122)	0.22	0.20	- 0.30	0.163	± 0.002	$^{+0.001}_{-0.003}$
$Scale \mu$	PARJ(129)	0.002	0.0005	- 0.0025	0.00250	± 0.00006	$^{+0.00013}_{-0.00018}$
a	PARJ(41)	0.5	0.1	- 0.5	0.903	± 0.014	$^{+0.038}_{-0.014}$
b	PARJ(42)	0.9	0.525		optimized		
σ_q	PARJ(21)	0.35	0.36	- 0.44	0.477	± 0.004	$^{+0.005}_{-0.006}$
$P(^1S_0)_{ud}$	-	0.5	0.3	- 0.5	0.327	± 0.023	$^{+0.051}_{-0.021}$
$P(^3S_1)_{ud}$	-	0.5	0.2	- 0.4	0.274	± 0.061	$^{+0.012}_{-0.013}$
$P(^1P_1)_{ud}$	-	0.	see text		0.089		
$P(oth.P - states)_{ud}$	-	0.	see text		0.310		
γ_s	PARJ(2)	0.30	0.27	- 0.31	0.277	± 0.006	$^{+0.008}_{-0.004}$
$P(^1S_0)_s$	-	0.4	0.3	- 0.5	0.413	± 0.028	$^{+0.016}_{-0.011}$
$P(^3S_1)_s$	-	0.6	0.2	- 0.4	0.330	± 0.021	$^{+0.011}_{-0.008}$
$P(P - states)_s$	-	0.	see text		0.257		
ϵ_c	PARJ(54)	-	-0.0950		optimized		
$P(^1S_0)_c$	-	0.25	0.26				
$P(^3S_1)_c$	-	0.75	0.44		adj. to data		
$P(P - states)_c$	-	0.	0.3				
ϵ_b	PARJ(55)	-	-0.00427		optimized		
$P(^1S_0)_b$	-	0.25	0.175				
$P(^3S_1)_b$	-	0.75	0.525		adj. to data		
$P(P - states)_b$	-	0.	0.3				
$P(qq)/P(q)$	PARJ(1)	0.1	0.08	- 0.11	0.087	± 0.002	$^{+0.008}_{-0.002}$
$P(us)/P(ud)/\gamma_s$	PARJ(3)	0.4	0.65		adj. to data		
$P(ud1)/P(ud0)$	PARJ(4)	0.05	0.07		adj. to data		
extra baryon supp.	PARJ(19)	0.	0.5		adj. to data only uds		
extra η supp.	PARJ(25)	1.0	0.65		0.65 ± 0.06		
extra η' supp.	PARJ(26)	1.0	0.23		0.23 ± 0.05		

Table 52: Parameter setting and fit results for JETSET 7.4 ME with default decays

F Tables of Correlation Coefficients

	a	σ_q	λ_{QCD}	Q_0	γ_s	$P(^1S_0)_{ud}$	$P(^3S_1)_{ud}$	$P(^1S_0)_s$	$P(^3S_1)_s$
a	1.00	0.33	-0.60	0.01	-0.10	0.27	-0.11	0.06	0.03
σ_q		1.00	-0.69	-0.08	0.43	-0.41	0.11	-0.26	-0.29
λ_{QCD}			1.00	0.18	-0.26	0.19	-0.10	0.15	0.20
Q_0				1.00	0.20	0.06	-0.38	-0.35	-0.13
γ_s					1.00	-0.04	0.19	-0.73	-0.65
$P(^1S_0)_{ud}$						1.00	0.37	-0.29	0.12
$P(^3S_1)_{ud}$							1.00	-0.10	-0.26
$P(^1S_0)_s$								1.00	0.41
$P(^3S_1)_s$									1.00

Table 53: Table of Correlation Coefficients for JETSET 7.4 PS Fit

	a	σ_q	λ_{QCD}	p_t^{QCD}	γ_s	$P(^1S_0)_{ud}$	$P(^3S_1)_{ud}$	$P(^1S_0)_s$	$P(^3S_1)_s$
a	1.00	0.11	-0.18	0.90	-0.29	0.71	-0.04	0.29	0.07
σ_q		1.00	-0.60	-0.07	0.12	-0.47	-0.06	0.02	0.05
λ_{QCD}			1.00	0.16	0.02	0.18	0.03	-0.03	-0.06
p_t^{QCD}				1.00	-0.21	0.65	-0.02	0.26	0.05
γ_s					1.00	-0.27	-0.02	-0.41	-0.47
$P(^1S_0)_{ud}$						1.00	-0.11	-0.03	0.13
$P(^3S_1)_{ud}$							1.00	0.08	-0.03
$P(^1S_0)_s$								1.00	0.02
$P(^3S_1)_s$									1.00

Table 54: Table of Correlation Coefficients for ARIADNE 4.06 PS Fit

G Figures of Data and Model Comparisons

The figures in this section compare DELPHI data and identified particle spectra to the following fragmentation models:

- JETSET 7.3 DELPHI decays labeled JT 7.3 PS
- JETSET 7.4 default decays labeled JT 7.4 PS
- ARIADNE 4.06 DELPHI decays labeled AR 4.06
- HERWIG 5.8 c default decays labeled H 5.8C
- JETSET 7.4 ME default decays labeled JT 7.4 ME

The references for the identified particle data can be found in table 7. For most shape distributions we show on the left side the data as measured from charged particles only, on the right as measured from charged plus neutral particles. The distributions are corrected to the corresponding final states. The total correction factor for each distribution is shown in the upper inset. The lower insets of the plots depict the relative deviation of the models to the data. Also shown as shaded area in these insets is the total experimental error obtained by adding quadratically the systematic and statistical error in each bin. Except for the point-to-point scatter the error should be interpreted like a statistical " 1σ " uncertainty.

G.1 Inclusive Distributions




Figure 3: Distribution of ξ_p and the scaled momentum x_p




Figure 4: Distribution of the Rapidity y_T and the Rapidity y_S




Figure 5: Distribution of p_t^{in} with respect to the Thrust axis




Figure 6: Distribution of p_t^{out} with respect to the Thrust axis




Figure 7: $\langle p_t^{out} \rangle$ vs. x_p and $\langle p_t \rangle$ vs. x_p




Figure 8: Distribution of the Energy Energy Correlation EEC and the Asymmetry AEEC

G.2 Shape Distributions




Figure 9: Distribution of 1-Thrust




Figure 10: Distribution of Major




Figure 11: Distribution of Minor




Figure 12: Distribution of the Oblateness




Figure 13: Distribution of the Differential 2-Jet Rate Durham Algorithm




Figure 14: Distribution of the Differential 2-Jet Rate Jade Algorithm




Figure 15: Distribution of the Differential 3-Jet Rate Durham Algorithm




Figure 16: Distribution of the Differential 3-Jet Rate Jade Algorithm




Figure 17: Distribution of the Differential 4-Jet Rate Durham Algorithm




Figure 18: Distribution of the Differential 4-Jet Rate Jade Algorithm




Figure 19: Distribution of the Sphericity




Figure 20: Distribution of the Aplanarity




Figure 21: Distribution of the Planarity




Figure 22: Distribution of the Difference of Jet masses




Figure 23: Distribution of the Heavy Jet mass




Figure 24: Distribution of the Light Jet mass




Figure 25: Distribution of the C-Parameter




Figure 26: Distribution of the D-Parameter




Figure 27: Distribution of the Wide Jet Broadening




Figure 28: Distribution of the Narrow Jet Broadening




Figure 29: Distribution of the Total Jet Broadening




Figure 30: Distribution of the Difference of the Jet Broadenings

G.3 Identified Particle Distributions




Figure 31: ξ_p Distribution of K^0




Figure 32: ξ_p Distribution of $K^\pm$




Figure 33: x_E Distribution of K^{*0}




Figure 34: x_E Distribution of $K^{*\pm}$




Figure 35: x_p Distribution of ρ and x_E Distribution of ϕ




Figure 36: x_p Distribution of f_0 and f_2




Figure 37: ξ_p Distribution proton




Figure 38: x_{i_p} Distribution Λ^0




Figure 39: x_p Distribution for Octet and Decuplet Baryons

References

- [1] DELPHI Coll., P. Aarnio et al., Nucl. Inst. Methods A303 (1991) 187
- [2] F. Jadach, B.F.L. Ward, Z. Was, Comput. Phys. Commun. 40 (1986) 285; Nucl. Phys. B253 (1985) 441
- [3] DELPHI Coll., DELPHI 89-67(1989); DELPHI 89-68(1989).
- [4] T. Sjöstrand, Comput. Phys. Commun. 39 (1986) 347; T. Sjöstrand and M. Bengtsson, Comput. Phys. Commun. 46 (1987) 367
- [5] J.E. Campagne and R. Zitoun, Z. Phys. C 43 (1989) 469
- [6] V. Blobel, Unfolding methods in high energy physics experiments, Lecture given at the 1984 School of Computing, DESY 84-118; V.B. Anikeev, V.P. Zhigunov, Phys. of Part. and Nucl. 24 Nr. 4 (1993) 424
- [7] H. Fürstenau, Ph. D. Thesis, IEKP-KA 92-16
- [8] L. Lönnblad, Comp. Phys. Comm. 71 (1992) 15.
- [9] G. Marchesini et al., Comp. Phys. Comm. 67 (1992) 465.
- [10] C. Peterson, D. Schlatter, I. Schmitt and P. Zerwas, Phys. Rev. D27 (1983) 105
- [11] B. Andersson, G. Gustafson, G. Ingelman, T. Sjöstrand, Phys. Rep. 97 (1983) 31.
- [12] G. Gustafson, U. Petterson, Nucl. Phys. B308 (1988) 746
- [13] F. Gutbrod, G. Kramer, G. Schierholz, Z. Phys. C21(1984) 235
K. Fabricius et al., Z. Phys C11(1981) 315
- [14] R.K. Ellis, D.A. Ross, E.A. Terrano, Phys. Rev. Lett. 45 (1980) 1225,
Nucl. Phys. B178(1981) 421
- [15] compare discussion and references in: QCD Generators, B. Bambah et al. CERN 89-08
Vol. 3 p. 143 ff.
- [16] P.M. Stevenson, Phys. Rev. D23 (1981) 2916
- [17] compare discussion and references in: QCD Generators, B. Bambah et al. CERN 89-08
Vol. 3 p. 235 ff.
- [18] ALEPH Coll., D. Busculic et al., Z. Phys. C 64, 361 (1994)
- [19] ALEPH Coll., D. Decamp et al., Z. Phys. C 55, 209 (1992)
- [20] TASSO Coll., M. Althoff et al., Z. Phys. C 26 (1984) 157; TASSO Coll., W. Braunschweig
et al. Z. Phys. C 41 (1988) 359
- [21] L3 Coll., B. Adeva et al., Z. Phys. C 55 (1992) 39
- [22] W.H. Press, B.P. Flannery, S.A. Teukolsky, W.T. Vetterling, Numerical Recipes, *The Art of Scientific Computing*, Cambridge University Press

- [23] NAG library MK16, NAG ltd., Oxford, GBR.
- [24] F. James, M. Goossens, *MIUNIT, Function Minimization and Error Analysis*, Reference Manual, CERN Program Library Long Writeup D506 (1992).
- [25] M.Z. Akrawy et al., Z. Phys. C 47 (1991) 505
- [26] V. Gibson, Charm and Beauty Hadron Production at $\sqrt{s} \approx M_Z$, Proc. XXVII ICHEP, Glasgow 1994, P.J. Bussey and I. G. Knowles (ed.)
- [27] ALEPH Collaboration, Production of K^0 and Λ in Hadronic Z Decays, CERN-PPE/94-74, submitted to Z. Phys. C.
- [28] OPAL Coll., R. Akers et al. CERN-PPE/94-206, submitted to Z. Phys. C.
- [29] DELPHI Coll., P. Abreu et al., Phys. Lett 345B (1995) 598
- [30] ALEPH Coll., D. Busculic et al., CERN-PPE/94-201, submitted to Z. Phys. C.
- [31] OPAL Coll., R. Akers et al., CERN-PPE/95-24 (1995), submitted to Z. Phys. C.
- [32] OPAL Coll., R. Akers et al., Z. Phys. C.63 (1994) 181
- [33] DELPHI Coll., P. Abreu et al., Nucl. Phys. B444 (1995) 3
- [34] OPAL Coll., Inclusive strange vector and tensor meson production in hadronic Z^0 decays, preliminary results presented at ICHEP94 Glasgow.
- [35] ALEPH Coll., preliminary results presented at ICHEP94 Glasgow, GLS0548. Holger Hepp (ALEPH Coll.),
Inklusive Produktion von geladenen K^* -Mesonen in hadronischen Z-Zerfällen, Diplomarbeit, HD-IHEP 93-04.
- [36] OPAL Collaboration, Strange Baryon Production and Correlations in Hadronic Z^0 Decays, preliminary results presented at ICHEP94 Glasgow.
- [37] DELPHI Coll., P. Abreu et al., Phys. Lett. 318B (1993) 249
- [38] DELPHI Collaboration, Strange Baryon Production in Z^0 Hadronic Decays. preliminary results presented at ICHEP94 Glasgow
- [39] DELPHI Coll., P. Abreu et al., Z. Phys. C65 (1995) 587
- [40] L3 Coll., Y. Pei, talk given at the CERN PPE Seminar
- [41] L3 Coll., O. Adriani et al., Phys. Lett. 286 B (1992) 403. L3 Coll., M. Acciari et al., Phys. Lett. 328 B (1994) 223
- [42] ALEPH Coll., D. Buskulic et al., Phys. Lett. 292 B (1992) 210. ALEPH Coll., Production Rates of η and η' in Hadronic Z Decays, preliminary results presented at ICHEP94 Glasgow.
- [43] DELPHI Coll., P. Abreu et al., CERN-PPE/95-39, submitted to Z. Phys. C.
- [44] Review of Particle Properties 1994, Phys. Rev. D50 (1994) 3

- [45] DELPHI Coll., P. Abreu et al., Z. Phys. C63 (1994)17, Phys. Lett. 323B (1994) 242, CERN-PPE/95-77, submitted to Phys. Lett.
- [46] A. De Angelis, J. Phys. G19, 1233 (1993)
- [47] OPAL Collaboration, M.Z. Akrawy Z. Phys. C 47 (1990) 505,
ALEPH Collaboration, D. Decamp et al., Phys. Lett. 273B(1991)181,
DELPHI Collaboration, A. Abreu et al., Z. Phys. C 50(1991)185,
OPAL Collaboration, P.D. Acton et al., Z. Phys. C 53 (1992) 539.
- [48] DELPHI Coll. P. Abreu et al., Phys. Lett B347 (1995) 447
- [49] Delphi Coll., P. Abreu et al. DELPHI 95-50 PHYS 488, contribution to the eps conference on High Energy Physics, Brussels 1995
- [50] S. Brandt et al., Phys. Lett. 12 (64) 57; E. Fahren, Phys. Rev. Lett. 39 (1977) 1587
- [51] J.D. Bjorken, S. Brodsky: Phys. Rev. D1 (1970) 1416
- [52] R.K. Ellis, D.A. Ross, A.E. Terrano: Nucl. Phys. B178 (1981) 421 G. Parisi: Phys. Lett 74B (1978) 65. J.F. Donohue, F.E. Low, S.Y. Pi: Phys. Rev. D20 (1979) 2759
- [53] L. Clavelli: Phys. Lett. B85 (1979) 111.
- [54] JADE Coll., W. Bartel et al.: Z. Phys. C33 (1986) 23 JADE Coll., S. Bethke et al.: Phys. Lett. B213 (1988) 235
- [55] L. Smolik, DESY 90-089
- [56] C.L. Basham, L.S. Brown, S.D. Ellis, S.T. Love: Phys. Rev. D17 (1978) 2298, Phys. Rev. Lett. 41 (1978) 1585, Phys. Rev. D19 (1979) 2018
- [57] S. Catani, G. Turnock, B.R. Webber, CERN-TH.6640/92 (1992)